

Publikasi 14

by Iyan 14

Submission date: 30-Nov-2017 01:17PM (UTC+0700)

Submission ID: 887517211

File name: Artikel_The_Effect_Of_Propylene_Glycol_Concentration_pdf.pdf (57.01K)

Word count: 2161

Character count: 11341

14 THE EFFECT OF PROPYLENE GLYCOL CONCENTRATION ON THE PHYSICAL CHARACTERISTIC AND RELEASE RATE OF CAFFEINE IN GEL

13 **Endang Diyah Ikasari, Milani**
Sekolah Tinggi Ilmu Farmasi "Yayasan Pharmasi" Semarang

Abstract

The aims of this research were to find if any the effects of the propylene glycol concentration on the physical characteristics of the gel which includes organoleptic, pH, viscosity, and of the caffeine releasing test within 360' that includes the value of dissolution-%, DE-%, lag time, and diffusion coefficient.

Caffeine gel formulation consisted of F I (with 10% of propylene glycol addition), F II (with 20% of propylene glycol addition), F III (with 30% of propylene glycol addition), IV (without of propylene glycol addition) and F V used as a positive control, an anti cellulite gel preparation that was available in the market used. Based on physical characteristic test, it could be drawn that the increasing propylene glycol concentration increased clarity, pH, and gel viscosity. Increasing propylene glycol concentration also decreased dissolution-%, decreased DE-%, increased lag time and decreased diffusion coefficient. The physical characteristic test and the caffeine releasing test were analyzed by using the one-way ANOVA with 95% and probability ($p=0,05$) level SPSS.

Based on the results of the research, it could be concluded that the different propylene glycol concentration as humectant, affects significantly to the physical characteristics, such as the pH and viscosity tests, and to the caffeine releasing in gel, such as the dissolution-%, DE-%, lag time, and diffusion coefficient.

Keywords : Caffeine, gel, propylene glycol, lag time, dissolution

INTRODUCTION

Cellulite is topographical skin alteration that happen in 85% post-adolescent women (Hexsel et al., 2006). It is normally occurred on the skin thighs, belly, and buttocks of whom that have an adrenergic receptor (Djajadisastra, 2008). Topical anticellulite is used to reducing cellulite. Caffeine is used as active ingredient that based on stimulation beta adrenergic receptor mechanism.

Hydrophylic gel are suitable to be used as cosmetic base because of its cooling effect, good appearance, and not sticky. Carbomer have also be chosen as gel-base because it can form a thicked gel (Allen, 2001) and produce a stable product (Amnuakit, 200). Gel component are usually consisted of humectant, a substance that can control water evaporation change between the product and the air, in the container or in the skin (Rukminingsih, 2009). Propylene glycol is common used as humectant that has mechanism in repairing irregular epidermis appearance of the skin cellulite condition. Propylene glycol can increase a solubility of many substances, one of which is caffeine (Rowe et al., 2009) and it can influence a swelling and viscoelasticity of carbomer (Islam et al., 2004). Propylene glycol is also known as a substance that influence release rate of active substances (Amnuakit et al., 2005).

METHODS

Object of this research was a physical characteristic and release rate of caffeine in gel as humectant in different concentration of 10%, 20%, and 30%. The parameter used to determine caffeine release rate was the value of lag time and coefficient diffusion based on Higuchi's Equation.

Free variable in this research was propylene glycol concentration of 10%, 20%, and 30%. Bound variable in this research were physical characteristics and release rate of caffeine. Physical characteristics gel consisted of organoleptic, value of pH, and viscosity. The value of lag time and diffusion coefficient could perform the rate of its caffeine's release.

The instruments used were mortar and stamper, glass tools, porcelain cup, *Hot Plate stirrer* model L-81 (Labinco), Tablet Dissolution Apparatus VDA-6DR *USP Standards* (VEEGO), pH 210 *Microprocessor* pH meter (HANNA), spektrofotometer UV mini 1240 *UV-Visible* (Shimadzu), viscosimeter Rion VT-04, and *paddle over disk*.

The materials used to form a caffeine gel, physical characteristic test and release rate were anhydrous caffeine, carbomer, propylene glycol, TEA, nipagin, KH₂PO₄, NaOH, cellophane membran, buffer pH 7 and buffer pH 4.

The method of this research was dissolving a caffeine into mixture of propylene glycol and water with magnetic stirrer until it dissolved well then added to carbomer gel base which was swelled and added enough water. Each formula consisted of a different propylene glycol concentration (table I).

Each formula has been tested its physical characteristic, of organoleptic, pH, and viscosity, and been done a release test of caffeine in gel using cellophane membrane and paddle over disk. The 2 grams of preparation gel were put into paddle over disk which cellophane membrane equipped, then put it into the dissolution vessel contained of 900 ml of phophat buffer pH 7,4 at 37°C, and was stirred at 50 rpm speeds. The sample was taken a 10,0 ml of it, at 5', 10', 15', 30', 45', 60', 90', 120', 150', 180', 240', 300', and 360' minutes stirring time. The absorbance of caffeine was viewed at the maximum wavelength of 273 nm on the 7,4 pH phosphate buffer blanko. The result datas were analyzed by using One way ANOVA for the normal distribution, and Kruskall Wallis for the abnormal distribution data.

Table I. The Calculation of component Caffeine Gel Preparation

Substances	Calculation	FI (g)	FII (g)	FIII (g)	FIV (g)
Caffeine	3% x 150 g	4,5	4,5	4,5	4,5
Carbomer	0,75 x 150 g	0,75	0,75	0,75	0,75
TEA	0,5% x 150 g	0,75	0,75	0,75	0,75
Propylene-Glycol	10% x 150 g	15	-	-	-
	20% x 150 g	-	30	-	-
	30% x 150 g	-	-	45	-
Nipagin	0,2% x 150 g	0,3	0,3	0,3	0,3
Aqua destilata	Hingga 150 g	Hingga 150	Hingga 150	Hingga 150	Hingga 150

RESULTS AND DISCUSSION


Based on the result on this research, it could be concluded that the increased propylene glycol concentration can increase clarity, pH's value, viscosity caffeine gel. Propylene glycol can increase caffeine solubility in gel and tend to be more alkalized, than carbomer that makes the carbomer swelling more easily, so that the gel clarity and pH were also increased. Carbomer swelling process causes increasing in the polymer cross-linked so then the viscosity is also increased. The result of the organoleptic test were shown at table II.

Table II. Result of Physical Characteristic and Caffeine Release Rate Test

Determine	Formula				
	F I	F II	F III	F IV	F V
Organoleptic:	Gel	Gel	Gel	Gel	Gel
Form	Clear	Clear	Clear	Turbid	Green and clear
Colour			(paling jernih)		
Odour	-	-	-		Tea's aromatic
Taste				-	
	Cool	Cool	Cool	Cool	Cool
pH	5,57 ± 0,03	5,75 ± 0,02	6,04 ± 0,02	5,20 ± 0,04	6,31 ± 0,02
Viscosity (P)	171,67 ± 2,58	182,50 ± 2,74	200 ± 0,00	158,33 ± 2,58	38,33 ± 5,16
Lag time (menit)	9,9987 ± 0,1092	16,8348 ± 0,5452	21,8593 ± 0,5473	7,2659 ± 0,2131	42,2219 ± 0,4138
Diffusion coefficient (10 ⁻⁷ x cm ² /minutes)	9,6967 ± 0,1470	5,7465 ± 0,1831	4,3986 ± 0,1093	13,2191 ± 0,3799	2,2763 ± 0,234
% Disolution	44,4888 ± 2,1111	35,3835 ± 1,3010	31,2257 ± 2,2495	54,8373 ± 3,7105	3,7876 ± 0,2986
% DE (Dissolution Efficiency)	56,2986 ± 3,7196	54,2677 ± 1,8589	51,5982 ± 2,1647	58,1339 ± 3,1156	52,7523 ± 1,8602

Caffeine release test was used to know the total caffeine solubilized and caffeine release rate. Caffeine release test was used negative control, that was a preparation gel without adding propylene glycol and the positive control, that was anti cellulite preparation that available in the market. Based on the result data, the relation between solubilized caffeine with in certain time interval, and the total solubilized caffeine of each formula (picture 1).

Formula I shoed a highest value of dissolution-% among other formulas, because amount of adding propylene glycol was just 10% and consisted more water than of formula II and formula III, so caffeine were more easily solubilized into the dissolution medium. The different water amount in each formula actually didn't give a significant effect on the caffeine gel preparation's release, if it was compared to the formula IV as negative control that showed the highest amount release because at the most of it preparation contained water, and without be added with propylene glycol. The negative control showed a highest release amount, because it did not contain a inhibitor of releasing factor such as propylene glycol. Formula V as positive control is showed a lowest result. It was because the preparation was formulated for slow release and consisted of other active anticellulite beside caffeine. Based on statistical test showed that increased propylene glycol concentration of 10% did not make any significant difference reduction of dissolution-%.



Picture 1. Caffeine dissolution content

Having in mind the levels of dissolution, the it was calculated the levels of Dissolution Efficiency (DE) that states the ratio between the area under the curve in the time t and the rectangular area that limits the 100% ordinate and absciss. Area under the curve or AUC describes the levels of drug content in time as a measure of size ant total of the bioavailability of drug. DE usulally used to determine the ability or efficiency of the drug is dissolved in the media until a certain time. Increased concentrations of 10% propylene glycol in the formula however did not cause a decrease in DE-% value significantly. While 20% increased propylene glycol concentration in the formula caused a decrease in DE-% value significantly.

The release rate of caffeine in gel were counted by applying the Higuchi equation. Caffeine release rate could be seen from the value of lag time and the value of diffusion coefficient. Release rate of active substance in gel were influenced by physical gel characteristic that was gel viscosity. Increasing gel viscosity was caused by density of polymer crossed-linked forms. Increasing the crossed-linked polymer density caused a gel hidrophobisity and inhibiting release rate of active substance (Sinko, 2006 : 519). A hidrophobisity also causes steric hydrance effect hydro so that inhibiting hydroxyl group interaction with its medium, that inhibits caffeine release rate.

Based on the result it concluded that, increasing propylene glycol leads to the increasing of lag time value and decrease the diffusion coefficient. The increased lag time value showed an inhibiting caffeine release rate from the gel basis up to the limit between its base and membrane surface. Based on the statistical test of lag time and diffusion coefficient, each formula tested with various concentration of propylene glycol showed significant differences.

Caffeine gel preparation as anticellulite worked locally and didn't enter systemic circulation. Adding propylene glycol as humectant in gel preparation are expected to improve skin appearance in cellulite condition and to restrain caffeine release rate in order to keep it work locally and doesn't enter the systemic circulation.

CONCLUSION

Based on the result of this research, can be concluded that :

1. Increased propylene glycol concentration gives significant effect to increase clarity, pH value, and viscosity value of caffeine gel preparation.
2. Increased 10% of propylene glycol concentration in the formula doesn't give significant effect of caffeine dissolution-% and DE-%.

3. Lag time value is increased and diffusion coefficient value is decreased, as result of increasing propylene glycol concentration.
4. The best formula is Formula III. It has the most clear physical characteristic, the longest release rate, and the less caffeine content, and best function as humectant.

REFERENCE

- 7 Allen, L.V. 2002. *The Art, Science and Technology of Pharmaceutical Compounding*. 2nd Ed. Washington DC : American Pharmacists Association
- 1 Amnuakit, T., Ikeuchi, I., Ogawara, K., Higaki, K., dan Himura, T. 2005. Skin Permeation of Propranolol from Polymeric Film Containing Terpene Enhancers for Transdermal Use, *Int J. Pharm* 289 (1-2) : 167
- 1 Amnuakit, T., Ingkawatornwong, S., Mancenuan, D., dan Worachotekamjorn, K. 2008. Caffeine Topical Gel Formulation, *IJPS* 4 (1) : 16-24.
- 9 Djajadisastra, J. 2008. *Effect of cream, gel and Ointment dosage forms on in-vitro skin penetration caffeine using franz diffusion cell*. UI repository. <http://repository.ui.ac.id/contents/koleksi/11/5a7469fb52aa316e1cffb9c5f18201e8204cb148.pdf> (28 Agustus 2010).
- 2 Hexsel, D., Zechmeister do Pado, D., Rao, J dan Goldman, M.P. 2006. Topical Management of Cellulite. In Goldman, M.P., Bacci, P.A., Leibaschoff, G., Hexsel, D., and Angelini, F (Eds). *Cellulite Pathophysiology and Treatment*. New York : Taylor and Francis Group.
- 5 Islam, M.T., Rodry guez-Hornedo, N., Ciotti, S dan Ackerman, C. 2004. Rheological Characterization of Topical Carbomer Gels Neutralized to Different pH, *AAPS* 21 : 7.
- Rowe, R.C., Sheskey, P.J dan Quinn, M. E. 2009. *Handbook of Pharmaceutical Excipient*. 6th Ed. Washington DC : Pharmaceutical Press and American Pharmacists Association
- Rukminingsih, F. 2009. Studi in vitro Pengaruh Gliserol, Propilenglikol, dan Sorbitol terhadap Pelepasan Perak Sulfadiazin dari Krim Setomakrogol. *Proceeding Seminar Nasional Farmasi "Antioksidan dalam Sediaan Obat, Kosmetika, Makanan dan Minuman"*. STIFAR "YAYASAN PHARMST" Semarang : 309-316.
- 8 Sinko, P. J. 2006. *Martin's Physical Pharmacy and Pharmaceutical Science*. 5th Ed. Philadelphia : Lippincott Williams & Wilkins.

Publikasi 14

ORIGINALITY REPORT

11%

SIMILARITY INDEX

9%

INTERNET SOURCES

9%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

mp.pharmacy.psu.ac.th

Internet Source

2%

2

www.ncbi.nlm.nih.gov

Internet Source

1%

3

Selvi Rahmawati, Riswi Haryatfrehni, Afra Meilianda, Matin Nuhamunada, Listia Pradani, Bagas Lantip Prakasa. "PIGMENTS CHARACTERIZATION OF MACROALGAE IN DRINI BEACH, GUNUNGKIDUL, YOGYAKARTA FOR SYSTEMATICS STUDY", KnE Life Sciences, 2015

Publication

1%

4

docplayer.net

Internet Source

1%

5

Submitted to University of Huddersfield

Student Paper

1%

6

eprints.ums.ac.id

Internet Source

1%

7

Submitted to University of Brighton

8

Submitted to University of Central Lancashire

Student Paper

1%

9

fhianunikoe.blogspot.com

Internet Source

<1%

10

id.scribd.com

Internet Source

<1%

11

www.dovepress.com

Internet Source

<1%

12

Kunihiko Gekko, Shozo Koga. "The stability of protein structure in aqueous propylene glycol", Biochimica et Biophysica Acta (BBA) - Protein Structure and Molecular Enzymology, 1984

Publication

<1%

13

ejournal-s1.undip.ac.id

Internet Source

<1%

14

Pramod P. Sarpotdar. "Evaluation of penetration enhancement of lidocaine by nonionic surfactants through hairless mouse skin in vitro", Journal of Pharmaceutical Sciences, 02/1986

Publication

<1%

15

Walker, J. C. F.. "Sawmilling", Primary Wood Processing, 1993.

<1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off