



Digital Receipt

This receipt acknowledges that **Turnitin** received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Maria Caecilia Nanny Setiawati
Assignment title: PENELITIAN
Submission title: Systemic Lupus Erythematosus Outpatients' Medication Kno...
File name: Revisi_65344-186283-1-RV_turnitin_1.docx
File size: 62.49K
Page count: 8
Word count: 2,842
Character count: 15,192
Submission date: 08-Mar-2023 12:38PM (UTC+0700)
Submission ID: 2031875215

JPSCR: Journal of Pharmaceutical Science and Clinical Research, 20ks
DOI: xxx/pscr.vxxx

Systemic Lupus Erythematosus Outpatients' Medication Knowledge and Quality of Life

Abstract

According to the Indonesian Lupus Foundation (YLI) in the last decade, the number of people with SLE (Systemic Lupus Erythematosus) has continued to increase every year. This research was undertaken to assess the medication knowledge and quality of life among SLE outpatients who had been counselled by pharmacist. A cross-sectional, non-experimental, descriptive study design was used in this research. A total of 121 respondents from Semarang and Yogyakarta met the criteria. Seventy (57.85%) respondents had previous counseling, while 51 (42.15%) respondents never. Their knowledge score was 32.17 ± 3.52 and the quality of life was 59.23 ± 17.04 . There was no correlation between the knowledge score and their quality of life ($P=0.014$). Their average age was 31.53 ± 7.77 years. It was positively related with their Quality of life ($P=0.031$) but not to their knowledge score. The participants' education level and disease duration were not considerably related with QoL and knowledge. Previous counseling by the pharmacist was positively related to their knowledge score, but not to their quality of life ($P=0.034$ and 0.793). The conclusion that can be drawn is that Pharmacists' counseling can improve SLE outpatients' knowledge scores significantly but not their QoL.

Keywords: SLE; outpatients; knowledge; QoL; pharmacist' counselling

1. Introduction

Systemic Lupus Erythematosus (SLE) is an autoimmune, inflammatory, chronic disorder characterized by the participation of the multigene system (Gomez et al., 2021). The disease impacts people worldwide, although the incidence and prevalence vary between countries. In Brazil, the prevalence ranges from 20 to 150 cases per 100,000 persons with the most prevalence were reported, and it appears to be growing more and more. People of Black/African American or Asian origin have a higher risk for SLE than White/Caucasian individuals, and show more serious disease phenotypes (Chen et al., 2022). Some data in Indonesia are obtained from hospitalized patients. In the year 2014, there were 1169 SLE in-patients but in 2016, there were 2166 in-patients diagnosed with SLE, in many hospitals in Indonesia, it was an 85% increased (Kementerian Kesehatan, 2017).

There are still few studies that show the knowledge of patients especially in chronic autoimmune cases (such as SLE) whose incidence is increasing nowadays and many people still don't understand all about SLE, cause nowadays SLE is still regarded as an uncommon disease in Indonesia. To characterize the full spectrum of the effects of SLE, Quality of Life is being known as an important aspect and relevant measure of efficacy in a clinical trial. Analyzing the

Systemic Lupus Erythematosus Outpatients' Medication Knowledge and Quality of Life

by Maria Caecilia Nanny Setiawati

Submission date: 08-Mar-2023 12:38PM (UTC+0700)

Submission ID: 2031875215

File name: Revisi_65344-186283-1-RV_turnitin_1.docx (62.49K)

Word count: 2842

Character count: 15192

Systemic Lupus Erythematosus Outpatients' Medication Knowledge and Quality of Life

Abstract

According to the Indonesian Lupus Foundation (YLI) in the last decade, the number of people with SLE (Systemic Lupus Erythematosus) has continued to increase every year. This research was undertaken to assess the medication knowledge and quality of life among SLE outpatients who had been counselled by pharmacist. A cross-sectional, non-experimental, descriptive study design was used in this research. A total of 121 respondents from Semarang and Yogyakarta met the criteria. Seventy (57.85%) respondents had previous counseling, while 51 (42.15%) respondents never. Their knowledge score was 32.17 ± 3.52 and the quality of life was 59.23 ± 17.04 . There was any correlation between the knowledge score and their quality of life. *P 0.014. Their average age was 31.53 ± 7.77 years. It was positively related with their Quality of life *P 0.031 but not to their knowledge score. The participants' education level and disease duration were not considerably related with QoL and knowledge. Previous counseling by the pharmacists was positively related to their knowledge score, but not to their quality of life (*P 0.034 and 0.793). The conclusion that can be drawn is that Pharmacists' counseling can improve SLE outpatients' knowledge scores significantly but not their QoL.

Keywords: SLE; outpatients; knowledge; QoL, pharmacist' counselling

17

1. Introduction

Systemic Lupus Erythematosus (SLE) is an autoimmune, inflammatory, chronic disorder characterized by the participation of the multiorgan system (Gomez et al., 2021). The disease impacts people worldwide, although the incidence and prevalence vary between countries. In Brazil, the prevalence ranges from 20 to 150 cases per 100,000 persons with the most prevalence were reported, and it appears to be growth more and more. People of Black/African American or Asian origin have a higher risk for SLE than White/Caucasian individuals, and show more serious disease phenotypes (Chen et al., 2022). Some data in Indonesia are obtained from hospitalized patients. In the year 2014, there were 1169 SLE in-patients but in 2016, there were 2166 in-patients diagnosed with SLE, in many hospitals in Indonesia, it was an 85% increased (Kementrian Kesehatan, 2017).

There are still few studies that show the knowledge of patients especially in chronic autoimmune cases (such as SLE) whose incidence is increasing nowadays and many people still don't understand all about SLE, cause nowadays SLE is still regarded as an uncommon disease in Indonesia. To characterize the full spectrum of the effects of SLE, Quality of Life is being known as an important aspect and relevant measure of efficacy in a clinical trial. Analyzing the

1 quality of life (QoL) is thus an important measure to examine how much the chronic disease process is influencing the SLE patients. There is a paucity of literature on QoL issues in SLE patients from Asia (Chen et al., 2022).

2 Pharmaceutical care needs a direct connection between a pharmacist and a patient as an individual. Patient counseling is one of the most important roles that a pharmacist can play. Patient's therapy accomplish counseling produce an occasion for pharmacists to enhance patients' therapeutic outcomes (Sanii Y et al, 2016). This study was conducted to know the sociodemographic profile of the SLE outpatients, to evaluate the knowledge about their medication and illness, and also to measure their quality of life. We also evaluate to know its correlation with age, disease duration, level of education, and being given previous counseling or formal information about SLE.

2. Material and Methods

We have obtained ethical clearance approval from the Ethics Committee of the Community Health Faculty Diponegoro University Semarang No 004/EC/FKM/2015 in a research to compile and developing Pharmacist counseling Module about SLE (This research article is a part of compiling that Module.)

Materials

We used questionnaire to measure the SLE outpatients' level of knowledge about SLE that has been validated before (Setiawati MCN, 2014). The questionnaire was prepared to measure the level of SLE outpatients knowledge about SLE. The questionnaire consisted of respondents' personal data (also whether the subjects has been given formally information) and 36 questions about understanding, causes and consequences of SLE (15 questions), non-pharmacological therapy (5 questions), SLE pharmacological therapy (9 questions), and drug's side effects (7 questions).

We also use questionnaires on Quality of Life (SF36) which has been validated and widely used all over the world. The SF36 is a generic, 36 item self-report questionnaire. It includes two component scores: the physical and the psychological component summary score. The SF-36 Score, a global measure of health-related quality of life, has been progressively reported in the scientific literature. Some studies using the SF-36 Score were published in a reputable journals (Liliane and Martins, 2016).

Methods

Non-experimental and a cross sectional study design was used in this research. Purposive samples were used to choose the subject, who met the criteria (had been diagnosed for more than

6 months, got routine therapy and agreed to be the subject of this research). Patients were notified of the objectives of the study and verbal consent obtained.

Respondents were asked to fill out a questionnaire (informed concern form, agreement form, and 36 questions) as a measure of their knowledge, and SF36 questionnaire as a measure of their quality of life.

Questionnaire, which was the instrument used for collecting information, was designed to obtain demographic data such as age, year of diagnosed, and level of education. This study was carried out in Middle Java: Semarang and Yogyakarta city. Data demography patients were shown descriptively and the other data was analyzed using SPSS 23. The inclusion criteria: patients have been diagnosed SLE (from physician, according to the American College of Rheumatology (ACR) for more than 6 months, got routine therapy and agreed to be the subject of this research (fulfilled all the questionnaires).

The filled questionnaires were analyzed as per the study objectives using Microsoft Excel. The SPSS version 23.0 was used to carry out the descriptive statistics. From the knowledge questionnaires, right answers were scored 1 and wrong answers were scored 0, and then we summarize the right answers as their level of knowledge, according to literature (Riwidikdo, 2013)

Score > mean +1,5 SD (> 35,95 → 36) as EXCELLENT category

Mean + 0,5 SD < score < mean +1,5 SD (33,93 < x < 35,95 → 34,35) as GOOD category

Mean -0,5 SD < score < mean + 0,5 SD 30,91 < x < 33,93 → 31,32,33) as ENOUGH category

Mean - 1,5 SD < score < mean-0,5 SD (26,39 < x < 30,91 → 27,28,29,30) as LESS category

Score < mean -0,5 SD (< 26,39 → 25,26) as POOR category

The SF 36 is the most valid generic health-related QoL questionnaire used in SLE (Kasitanon et al, 2013). Bearing in mind that QoL is best measured by the SLE patients himself/herself and not his or her physician or nurse or other people [Slevin et al, 1988], the questionnaire was administered to the patients. Domain scores were calculated according to the SF36 guidelines. The more the score is representing better QoL. We divided the QoL score into 5 groups, 60 or more as a GOOD category and less than 60 as a BAD category

Disease duration was described as the time from when diagnosis of SLE was made by a physician. Predictive factors in this research included age, level of education, disease duration and previous counseling. We determined if there is any correlation of SLE outpatients' level of knowledge and their quality of life, with their level of education, disease durations and ever being counseled before. Interrelation between variables was analyzed statistically using the cross-tabulation (Chi square) method implemented in SPSS v23. A $P < 0.05$ was considered significant.

Commented [A1]: Penulisan sitasi tidak tepat

Commented [A2R1]: Sudah menggunakan Zotero type JPSCR. Penulisnya 3 orang. Tks

Commented [A3]: Analisis datanya dari total skor kuesioner apa dikategorikan dulu. Akan lebih terlihat hubungannya kalau dari total skor. Analisisnya bisa pakai Pearson atau Spearman ttt normalitas data.

Commented [A4R3]: Data asli dari skor hasil pengisian kuesioner. Tks. Analisa menggunakan chi square. Tks

3. Results and Discussion

One hundred and thirty-six SLE outpatients were approached and accepted to participate in the study, only 121 participants completed the data and fulfilled the questionnaires, giving a response rate of 88.97%. Among them, 1.65 % (n=2) were males and 98.35 % (n=119) were females. We don't compare the knowledge and the quality of life between them, because the male amount was so little and not comparable.

The prevalence of SLE is consistently reported to be higher in females than in males, with a ratio of approximately 10:1 (Waldheim Eva, 2013). Although SLE can affect both men and women at any age, 90% of individuals diagnosed with the disease are female, and 80% of all patients with systemic lupus progress it between ages 15 and 45 (Howard and Mayhew, 2006). Another study found that about 80–90% of all systemic lupus erythematosus cases are women in the age of 20 to 40 years (with mean age at diagnosis: 29 years) (Olesinska and Saletra, 2018). The average age was 31.53 ± 7.77 years, the youngest is 16 years old and the oldest is 56 years old. This was largely similar to other studies, cause SLE is dominant in women in productive age (Shahin D, 2011). In this study, the SLE patients' disease duration is 5.80 ± 5.0 years. The shortest has been diagnosed only six months before the research, the longest has been diagnosed for 33 years by the clinician. This shown that SLE is a chronic disease, which stated in many studies (Chen et al., 2022). The demographic data of the respondents are shown in table 1.

Table 1. Demographic characteristic of survey respondents (n=121)

Variables	N	%	range	mean $\pm$ SD
Gender: Male	2	1.65		
Female	119	98.35		
Age (years)	121		16-56	31.53 ± 7.77
< 30 years	52	42.97		
30-40 years	46	38.02		
>40 years	23	19.01		
Disease duration (years)	121		0.5-33	5.80 ± 5.0
< 5 years (BF)	66	54.55		
≥ 5 years (AF)	55	45.45		
Education				
below high school	5	4.13		

Commented [A5]: Berdasarkan jenis kelamin. Supaya ditambahkan di atasnya

Commented [A6R5]: Sudah kami tambahkan: Gender. Tks

Commented [A7]: Yang bertanda kuning ini agak membingungkan pembaca. Sebaiknya dinotasikan saja.

Commented [A8R7]: Narasi saya tambahkan di atas tabel 1. Tks

high school	50	41.32
graduate	66	54.55

The cross-sectional design in this study, as well as the relatively small sample, may limit the generalization of the results. However, despite the sample size, several significances were found. On the other hand, because of the sample size, further significances may have been undetected.

Of all the respondents, the mean knowledge score was 32.17 ± 2.52 (enough category), Two (1.65%) respondents got the lowest score, namely 25, and 11 (9.09%) respondents got the highest score, 36. The mean of their quality of life score was 59.23 ± 17.04 (bad category). There was a correlation between the knowledge score and the quality of life, with $*P 0.014$. The higher the knowledge scores, the better the Quality of life. The level of the subjects' knowledge about SLE can be seen in figure 1.




Figure 1. The subject's knowledge about SLE

The association between the age of patients, level of education, disease duration and ever being counseled before with knowledge and Quality of Life score was analyzed and the results were summarized in Table 2. There was no serious interrelation between the age of the patients and their knowledge $P > 0.05$. However, there was a relationship between age and patients' quality of life, $*P 0.031$. The level of education of the SLE outpatients did not have any correlation with their knowledge score and quality of life. This result, different from other research (Alkatheri and Albekairy, 2013) which found the significant interrelation between the level of education and the patient knowledge.

Commented [A9]: Sebaiknya dibuat tabel

Commented [A10R9]: Done, figure 1. tks

Commented [A11]: Bisa ditambahkan korelasinya seperti apa. Misalnya makin tinggi skor pengetahuan, makin baik kualitas hidupnya.

Commented [A12R11]: Done, tks

Many studies showed enhancements in patient knowledge of their medication as a result of pharmacist counselling interventions (Schmitt MR et al, 2011). The need for pharmacist's counseling, is needed by patients, especially for SLE patients whose disease is not yet well known and the treatment varies greatly. Our data demonstrated that Indonesian SLE outpatients' knowledge was not satisfied and had relatively poor quality of life.

The mean of the disease duration was 5.80 ± 5.0 year. study the influence of disease duration, the study population was divided into two groups based upon the median disease duration: those with a disease duration below 5 years were 66 patients (54.55%) and equal or more than 5 years were 55 patients (45.45%). There was no such a relationship between the disease duration of the patients with their knowledge and Quality of life, $P = 0.061$ and 0.054 .

The participants were also categorized according to their level of education into three categories: elementary school (below high school): only 5 patients (4.13 %), high school: 50 patients (41.32%) and graduated: 66 patients (54.55 %). Their knowledge mean were 41.5, 63.75 and 60.39 and their QoL mean were 51.1, 58.13 and 63.92. There was no correlation between level education with their knowledge and Quality of life ($P = 0.386$ and 0.551). There was no such a relationship between the patients' education with their medicine knowledge and Quality of life, $P = 0.371$ and 0.773 . The Quality of Life did not have any relationship with disease duration, level of education and ever received previous counseling, $P > 0.05$.

Finally, 70 patients (57.85 %) of 121 participants had received previous information about SLE as a counseling from the physicians but not from the pharmacist, and the other 51 patients (42.15%) of the patients did not receive previous counseling or they cannot remember, respectively. There was no significant association between the patients' ever received previous counselling with their Quality of life, $P > 0.05$. However, the patients' knowledge was found to be significantly associated with ever received previous counselling, $P < 0.05$.

The quality of life (QoL) is being recognized as an important aspect of chronic disease. QoL is difficult to measure and there is no consensus on the single best instrument to use. Most researches report a faulty QoL in lupus patients (Wang et al, 2001). Health-related quality of life is an independent territory in the analysis of patients with SLE, and actually the QoL is lower in patients with SLE than in patients with other chronic diseases (Kuriya B et al, 2008). Another study also made a conclusion that the quality of life in patients with SLE is lower than in general population and comparable with other chronic diseases (da Hora et al., 2019).

The QoL of SLE patients decreases time after time. Using the SF36, several studies have demonstrated that persons with SLE have a significantly poorer QoL than persons without a

Commented [A13]: Menggunakan uji apa?

Commented [A14R13]: Association between variables was statistically analyzed using the cross-tabulation (Chi square) method implemented in SPSS v23. A $P < 0.05$ was considered significant. Sudah tercantum pada Metode, tks

Commented [A15]: Paragraf tidak boleh hanya satu kalimat

Commented [A16R15]: Saya gabungkan dengan paragraph diatasnya. Tks

chronic illness (Morales E et al. 2017). In this study, the patients' quality of life was evaluated substantially related with age, P 0.031. The other study, stated that age has a negative influence on QoL, especially physical health (Mc Elhone, 2006). In our study, the disease duration was not associated with quality of life, P 0.054. Another study stated that the effect of disease duration on Quality of life was still unclear, and cause of the confirmed and inveterate of SLE, it will need repeated assessment of health-related quality of life (HRQoL) (Chaigne et al., 2017). There are several studies reporting mutual relationships between subjective symptoms in SLE, like pain, fatigue, anxiety and depression, as well as HRQoL (Pettersson et al, 2012). Another study, like our, found that Educational level and disease duration were not associated with HRQOL domains (Etchegaray Morales, 2017). In general, the results from SF-36 are in line with previous studies in other populations showing impaired health-related quality of life in patients with SLE differentiated from controls (McElhone et al. 2006).

Table 2: The QoL and Knowledge of SLE' outpatients (n=121)

Variables	N	%	mean $\pm$ SD
Previous counseling/PC			
No	51	42.15	
yes	70	57.85	
QoL	121		59.23 $\pm$ 17.04
BF	66		56.86 $\pm$ 2.22
AF	55		61.92 $\pm$ 2.11
No PC	51		58.89 $\pm$ 16.85
Yes PC	70		59.36 $\pm$ 17.38
Knowledge	121		32.17 $\pm$ 2.52
BF	66		33.34 $\pm$ 2.11
AF	55		31.99 $\pm$ 2.7
No PC	51		31.77 $\pm$ 2.6
yes PC	70		32.73 $\pm$ 2.32

Fifty-one patients (42.15 %) had previous counseling before but non of them being counselled by pharmacist, they were counselled by physician and the other by survival SLE patients. There was significantly associated between Previous counseling with knowledge (p

0.039). Many researches have shown that counselling achieved by a clinical pharmacist upgrades patients' knowledge of their medications (Ganachari and Almas, 2012). Education and knowledge of patients to appropriate use of medicines and afterward positive therapeutic outcomes and disease control could motivate patients in medication compliance (Sanii Y et al, 2016).

4. Conclusion

The most important finding of the present study was many of the SLE outpatients had low knowledge and that they need special information about their disease and especially about their therapy. SLE outpatients' quality of life is still not good. Providing information such as counseling by pharmacist can improve SLE outpatients' knowledge but not their quality of life. Level of education and disease duration did not affect the QoL in any of the domains.

Acknowledgement

At first, we would like to warmly thank the SLE outpatients who participated in this study and who generously gave of their time to be the subjects of this research

The author wish to acknowledge:

Directorate of Research and Community Service

Directorate General of Research and Development Strengthening Ministry of Research Technology and Higher Education (RISTEKDIKTI) According Research Contract

23

Conflict of Interest

As the authors, we have declared there is no conflicts of interest.

References

Systemic Lupus Erythematosus Outpatients' Medication Knowledge and Quality of Life

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

13%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	rheumatology.oxfordjournals.org Internet Source	4%
2	www.ncbi.nlm.nih.gov Internet Source	1%
3	www.termedia.pl Internet Source	1%
4	Pantelis Panopalis, Ann E. Clarke. "Quality of Life in Systemic Lupus Erythematosus", Clinical and Developmental Immunology, 2006 Publication	1%
5	Bindee Kuriya. "Quality of life over time in patients with systemic lupus erythematosus", Arthritis & Rheumatism, 02/15/2008 Publication	1%
6	www.uspharmacist.com Internet Source	1%
7	Submitted to Brookdale Community College Student Paper	1%

8

LEILA F. DANTAS, IGOR T. PERES, LEONARDO S.L. BASTOS, JANAINA F. MARCHESI et al.
"App-based symptom tracking to optimize SARS-CoV-2 testing strategy using machine learning", Cold Spring Harbor Laboratory, 2020

Publication

1 %

9

www.researchgate.net

Internet Source

1 %

10

www.researchsquare.com

Internet Source

1 %

11

Submitted to University of Derby

Student Paper

1 %

12

Aya Ahmed, Mohammed Ismail, Sahar Elsayed, Rabab Hussein. "Quality of Life in Patients with Systemic Lupus Erythematosus", Sohag Medical Journal, 2018

Publication

1 %

13

Matthew W. Yung, Jim Gould, Graham J. G. Upton. "Nasal Polyposis in Children with Cystic Fibrosis: A Long-Term Follow-up Study", Annals of Otology, Rhinology & Laryngology, 2016

Publication

1 %

14

jurnal.uns.ac.id

Internet Source

1 %

15	Alaa Mohamed E.Isayed, Seham Guirguis Ragheb, Mona Abo Bakr. "Quality of Life among Clients with Benign Prostatic Hyperplasia", Egyptian Journal of Health Care, 2018 Publication	<1 %
16	Leopoldo Magacho, Francisco Lima, Ana Carla Nery, Antônio Sagawa, Bernardo Magacho, Marcos Ávila. "Quality of life in glaucoma patients: Regression analysis and correlation with possible modifiers", Ophthalmic Epidemiology, 2009 Publication	<1 %
17	cyberleninka.org Internet Source	<1 %
18	espace.library.uq.edu.au Internet Source	<1 %
19	Submitted to Grand Canyon University Student Paper	<1 %
20	talenta.usu.ac.id Internet Source	<1 %
21	core.ac.uk Internet Source	<1 %
22	Submitted to International Medical University Student Paper	<1 %

23 Mitra Tarlan, Sedigheh Khazaei, Seyed Hamid Madani, Elaheh Saleh. "Prognostic factors for cancer - specific survival in 220 patients with breast cancer: A single center experience", Cancer Reports, 2022
Publication

24 D. Spurgeon. "Children with asthma and their families benefit from asthma education", BMJ, 08/18/2009
Publication

25 Liliane Lins, Fernando Martins Carvalho. "SF-36 total score as a single measure of health-related quality of life: Scoping review", SAGE Open Medicine, 2016
Publication

26 S. Khanna, H. Pal, R. M. Pandey, R. Handa. "The relationship between disease activity and quality of life in systemic lupus erythematosus", Rheumatology, 2004
Publication

Exclude quotes Off
Exclude bibliography Off

Exclude matches Off