

Anti-inflammatory Activity of Salicylanilide Compounds from Gondopuro Oil as an In-Vitro

by Erwin Indriyanti

Submission date: 26-Sep-2024 01:31PM (UTC+0700)

Submission ID: 2465993090

File name: 1_Surat_Tugas_dan_Artikel_Salisilanilid_november_2023.pdf (1.41M)

Word count: 3402

Character count: 18197



SEKOLAH TINGGI ILMU FARMASI YAYASAN PHARMASI SEMARANG

LEMBAGA PENELITIAN DAN PENGABDIAN MASYARAKAT

Jalan Letnan Jendral Sarwo Edie Wibowo Km. 1 Plamongansari - Pucanggading - Semarang - 50193

Telepon : 024 - 6706147 ; 6725272 ; Faksimile : 024 - 6706148

Email : stifar_yaphar@yahoo.com

Website : www.stifar.ac.id

SURAT TUGAS

No. 068e/LK-TM/LPPM/STP/XI/2023

Yang bertanda tangan dibawah ini :

Nama : Dr. Lia Kusmita, S.Si.,M.Si

NIY : 040309003

Jabatan : Ketua LPPM STIFAR YAYASAN PHARMASI SEMARANG

Memberi tugas kepada :

No.	Nama	NIY	Jabatan
1.	apt. Dhimas Adhityasmara, M.Farm.	YP. 041015069	Dosen
2.	Dra. apt. Eka Susanti Hanhadyanaputri, M.Farm.	YP. 020991001	Dosen
3.	Erwin Indriyanti, S.Si., M.Pd.	YP. 040217099	Dosen
4.	apt. Aries Koes Sundoro, M.Farm.	YP. 040917113	Dosen

Untuk melaksanakan publikasi artikel ilmiah di **Jurnal Al-Kimia, Vol. 11; No. 2**, dengan judul "*Anti-inflammatory Activity of Salicylanilide Compounds from Gondopuro Oil as an In-Vitro*". pada :

Periode : Desember 2023

Tempat : UIN Alauddin Makassar

Demikian surat tugas dibuat kepada yang bersangkutan untuk dapat dilaksanakan dengan penuh rasa tanggung jawab.

Semarang, 18 November 2023

Ketua LPPM STIFAR

YAYASAN PHARMASI SEMARANG



[Signature]
Dr. Lia Kusmita, S.Si.,M.Si

NIY. YP. 040309003

Anti-inflammatory Activity of Salicylanilide Compounds from Gondopuro Oil as an In-Vitro

Dhimas Adhityasmara¹, Eka Susanti H. P.², Erwin Indriyanti³, Aries Koes Sundoro^{4*}

^{1,2,3,4} Department of Pharmacy, Stifar Foundation Pharmacy Semarang
Jl. Sarwo Edie Wibowo Km.1 Kota Semarang, Indonesia

*Corresponding Author: lutaris101010@gmail.com

Abstract: A component of gondopuro oil is methyl salicylate. Methyl salicylate has functional groups that allow it to be changed. Changes in functional groups can occur through chemical reactions, one of which is through the aminolysis reaction to produce salicylamide or salicylanilide. The precursor for the synthesis of amide derivatives is a carboxylic acid, using natural methyl salicylate from gondopuro oil and amines sonochemically using sonochemical methods for 3, 4, and 5 hours with temperature control of $\pm 60^{\circ}\text{C}$. Then liquid extraction was carried out using of hexane and distilled water 2 to 3 times until 2 phases were formed, then left for 1 night at a temperature of less than 10°C and decanted. Then extracted with 10 mL of cold 5% NaOH 2 times until two phases were formed, namely the n-hexane fraction and the NaOH fraction. The NaOH fraction obtained is then heated using a water bath to evaporate the solvent so that it becomes a solid, which is then placed in an oven to form crystals. The crystals obtained were then dried using an oven at 40°C until dry, and then the yield of the synthesized powder was calculated. From the research results, it can be concluded that Salicylanilide compounds can be synthesized using sonochemical methods and produce the largest % yield in the 3rd sample at 5 hours at 11.23%. Salicylanilide compounds have anti-inflammatory activity at a concentration of 100.0 ppm with a % inhibition of 46.07%.

Key word: Anti-inflammatory, Gondopuro oil, Inhibition, Salicylanilide, Sonochemistry

INTRODUCTION

Gondopura oil is a class of essential oils that is widely known by the general public. The main component of gondopuro oil is 96-99% methyl salicylate (Ma'mun 2015)(Sulistyo, Suratmo, and Retnowati 2015). This compound is widely used in the pharmaceutical industry, fragrances, and the food and beverage industry. Apart from that, the methyl salicylate compound has many properties so that it can be used as an anti-inflammatory, acute rheumatic medicine to relieve pain and relieve breathing, anti-inflammatory and analgesic (F. Dani Hendrata 2014) (Ripa, Dash, and Faruk 2015)(Of et al. 2015).

Methyl salicylate (o-hydroxymethylbenzoate) has functional groups that allow it to be converted into other compounds derived from methyl salicylate through changes in functional groups through organic chemical reactions. Changes in functional groups can occur through chemical reactions, including through the hydrolysis reaction to produce salicylic acid, through the aminolysis reaction to produce salicylamide or salicylanilide, through the transesterification reaction to produce other ester compounds such as ethyl salicylate, isoamyl salicylate, phenyl salicylate (salol) and etc (Hendrata 2014) (Retnowati and Azzuhro 2013). Other reactions that can change the functional groups in methyl

salicylate include alkylation, reduction, electrophilic substitution on the benzene ring and others (Retnowati and Azzuhro 2013).

The sonochemical method is used as an alternative to conventional stirring with low energy and efficient results (Draye, Chatel, and Duwald 2020) (Indriyanti and Prahasiwi 2020). This method is also effective for synthesizing compounds in a relatively short time. This research also uses a Decyclocarbodiimide (DCC) coupling reagent to speed up the coupling process of a reaction (Farshori et al. 2010).

Based on the above background, the researcher intends to synthesize amide derivatives with a sonochemical method times of 5 hours. Tests for the purity of synthetic compounds include melting point tests, solubility tests and thin-layer chromatography tests. Identification of synthetic compounds by spectrum elucidation using infrared spectrophotometry (FTIR-ATR), GCMS and in vitro anti-inflammatory activity testing.

RESEARCH METHODS

Materials and Tools

The ingredients used in this research were gondopuro oil, aniline (p.a), ethanol (p.a), distilled water, methanol (p.a), Bovine serum albumin (BSA). The tools used in this research are glassware in the laboratory, sonicator, IR spectrometer, spectrophotometer, melting point apparatus, GC-MS, vacuum rotary evaporator, Buchner filter, TLC plate, UV spectrophotometer -Vis, melting point apparatus and capillary tube.

Synthesis of Salicylanilides

12 mL of gondopuro oil was reacted with 9 mL of aniline, 50 mL of hexane and 10 mL of NaOMe 10% sonicated for 3, 4, and 5 hours at a temperature of 60°C. Then, liquid extraction was carried out using 50 mL of hexane and distilled water 2 to 3 times until 2 phases were formed, which is then separated in a separating funnel into a hexane fraction and a water fraction. The n-hexane fraction was added with anhydrous Na₂SO₄, then left for 1 night at a temperature of less than 10°C and decanted. Then extracted with 10 mL of cold 5% NaOH 2 times until two phases are formed, namely the n hexane fraction and the NaOH fraction. The NaOH fraction obtained is then heated using a water bath to evaporate the solvent so that it becomes a solid, which is then placed in an oven to form crystals. The crystals obtained were then dried using an oven at 40°C until dry, and then the yield of the synthesized powder was calculated (Sulistyo et al. 2015) (Sondhi et al. 2009).

Structure elucidation with Infrared Spectroscopy (FTIR-ATR) and GC-MS

The synthesized compound was placed on an ATR crystal and its spectrum was measured at a wave number of 4000-500 cm⁻¹ and structure elucidation with GC-MS

Anti-Inflammatory Test by In-vitro Activities

In vitro anti-inflammatory test based on the research method by (Williams et al., 2008). Stages testing activity of synthesis results in the denaturation of Bovine Serum Albumin (BSA). A total of 5 ml of positive control solution consisted of 4.950 l BSA and 50 l diclofenac sodium solution. The control solution was made in various concentrations, namely 100 ppm, 10 ppm, 1 ppm, and 0.1 ppm. Each of the solutions was vortexed and then incubated for 30 minutes at room temperature (27°C). After that, it was heated for 5 minutes at 72°C. Then, it was left at room temperature (27°C) for 25 minutes. Then the turbidity was measured using a UV-Vis spectrophotometer at a wavelength of 660 nm (Almira et al. 2021). The percentage of inhibition of BSA denaturation can be counted with this formula.

$$\%inhibition = \frac{Abs\ Negative\ Control - Abs\ Sample}{Abs\ Negative\ Control} \times 100\%$$

2

RESULTS AND DISCUSSION

The synthesis of salicylanilide compounds in this study was carried out based on the research method by (Retnowati and Azzuhro 2013) with slight modifications sonochemical methods. The resulting compound was then tested for FTIR-ATR and GCMS and potential as an anti-inflammatory activity in vitro. The physical properties of Gondopuro oil were determined and compared with the standard material safety data sheet (MSDS) for wintergreen oil shown in Table 1.

Table 1. Physical Properties of Gondopuro Oil

Physical Properties Parameters	Gondopuro Oil	Standard wintergreen oil (MSDS)
Being	Liquid	Liquid
Colour	Pale yellow	Colourless to yellowish
Smell	Sharp	-
Specific gravity	1.185 g/mL	1.18 g/mL

2

Based on Table 1, it can be seen that the form, color, and specific gravity of the gondopuro oil used in this research are close to the standard data contained in the material safety data sheet (MSDS) for wintergreen oil. Gondopuro oil was identified using FTIR to determine functional groups and GC-MS to determine the chemical structure and abundance of synthesized compounds. This analysis can also be used as a comparison for the results of FTIR and GC-MS analysis of reaction compounds.




Figure 1. FTIR Spectra Graph of Gondopuro Oil

The results of the spectrum analysis of gondopuro oil using FTIR in Figure 1 show that there is vibration absorption in the 3185 cm^{-1} area, namely stretching vibration =C-H sp^2 (aromatic ring) which is supported by absorption in the 753 cm^{-1} area, namely bending

vibration =C-H. The absorption in the 1674 cm^{-1} area is the C=O stretching vibration for esters. The functional group is supported by strong absorption in the 1301 cm^{-1} and 1251 cm^{-1} areas, namely C-O (ester) stretching vibrations which are shown by 2 peaks between $1300\text{--}1000\text{ cm}^{-1}$, then absorption appears in the 1584 cm^{-1} area, namely vibration stretch C=C for the aromatic ring. The absorption in the 1210 cm^{-1} area is the C-O stretching vibration for alcohol.

Identification using GC showed a peak at a retention time of 6,103 minutes with an abundance of 99.97% showing a mass spectrum with a molecular weight of 152 which indicated that the compound was a methyl salicylate compound. The GC-MS test results for gondopuro oil are shown in Figures 2 and 3.




Figure 2. Chromatogram of Gondopuro Oil




Figure 3. Mass spectrum of Gondopuro oil with a retention time of 6,100 minutes

Mass spectra results from GC-MS testing of Gondopuro oil showed that the fragmentation pattern of one of the components of Gondopuro oil included $m/z = 121$ due to the loss of the CH_3O radical ion (M-31), followed by the release of the CO carbonyl radical ion (M-31-28) producing a peak. $m/z = 93$. Peak $m/z = 93$ produces peak $m/z = 64$ (M-29) due to the loss of CHO radical ions. The peak $m/z = 51$ (M-13) occurs from $m/z = 65$, which loses CH radical ions. The fragmentation pattern of the methyl salicylate compound can be seen in Figure 4.




Figure 4. Fragmentation Pattern of Methyl Salicylate Compound

Salicylanilide can be synthesized directly by reacting methyl salicylate obtained from gondopuro oil with aniline solvent. Aniline is a nucleophile so the N atom in aniline can attack the C carbonyl atom which is double bonded to the O atom in methyl salicylate. This process causes a transitional attachment between aniline and methyl salicylate, where the N: atom of aniline will experience attachment to the C atom in methyl salicylate and the $-OCH_3$ group will experience a bond cleavage from methyl salicylate. The process is then continued with the release of the H atom which is bound to the N atom in aniline as the primary amine and forms an amide group, thus obtaining the salicylanilide compound. The synthesis of salicylanilide was carried out via a nucleophilic substitution reaction using the sonication method. If there is excessive activation of highly reactive intermediate compounds, it will cause intramolecular reactions to occur in the intermediate compound itself. Is intended to shorten the synthesis time where ultrasonic wave radiation can speed up the reaction. Ultrasonic waves when in a liquid medium can cause acoustic cavitation (Manickam et al. 2023), during the cavitation process bubble collapse will occur (bubble instability), namely the breaking of small bubbles due to sound. The synthesis results can be seen in Table 4.

Table 4. % Yield from Synthesis Results

Sample	Theoretical Weight (g)	Weight of Synthesis Results (g)	% yield
Sample 1 (3 Hour)	1.4841	0.1058	7.13
Sample 2 (4 Hour)	1.4841	0.1316	8.87
Sample 3 (5 Hour)	1.4841	0.1712	11.53




Figure 5. FTIR Spectra Graph of Reaction Result Compounds

Table 5. Data on Wave Numbers of Reaction Result Compounds

Number	Group	Reaction Result Compounds			References (Suratmo dkk., 2013)
		Sample 1	Sample 2	Sample 3	
1	O-H (phenol)	3347	3386	3390	3406.05
2	C-H aliphatic	3161	3051	3176	2991.39
3	C=O (Amide)	1674	1627	1588	1631.67
4	C=C aromatic	1433	1435	1444	1458
5	C-N aliphatic	1316	1258	1316	1253.64

Based on the wave number data in Table 5 and the FT-IR spectrum in Figure 5, it shows that there is absorption in the 3390 cm^{-1} area, namely the O-H stretching vibration of phenol, which is supported by absorption in the 1444 cm^{-1} area, namely the aromatic C=C stretching vibration. Next, absorption appears in the 3051 cm^{-1} area, which is the aliphatic C-H stretching vibration. The absorption for C=O (amide) is shown in the 1674 cm^{-1} area, while the absorption in the 1258 cm^{-1} area is the aliphatic C-N stretching vibration. Amides have a typical absorption in the 1680-1630 cm^{-1} region, namely C=O stretching vibrations. The results of the FTIR test of the reaction compound show that the wave number of the synthesized compound is close to the wave number in the research of (Retnowati and Azzuhro 2013).

The resulting compound was tested for anti-inflammatory activity using Bovine Serum Albumin (BSA). Protein denaturation is a process where proteins lose their tertiary and secondary structures by external compounds such as strong acids, strong bases, organic salts, organic solvents and heating. Generally, proteins lose their biological function when they are denatured. According to (Whittaker and Vogler 2008), compounds that have a percentage of protein denaturation inhibition of >20% are compounds that have anti-inflammatory activity. Testing of anti-inflammatory activity on protein denaturation was carried out by adding BSA solution, this was to reduce the use of live specimens in the drug development process and when BSA was heated, BSA denaturation would occur.

According to (Nasution, Nurilmala, and Abdullah 2019), compounds that can stabilize proteins from the protein denaturation process are compounds that have the potential to act as anti-inflammatories where there is an interaction between BSA and the active substance of the compound which results in the binding of the active substance with tyrosine, threonine and lysine. When the active substance sticks to the active substance it will not prevent BSA denaturation. The anti-inflammatory test using protein denaturation was carried out on the synthesized compound and Diclofenac Sodium as a positive control.

Table 6. In Vitro Anti-Inflammatory Test Results

Number	Sample	Concentration (ppm)	Absorbance	% Inhibition
1	Negative Control	-	0.280	-
		0.1	0.210	25.00
2	Na diklofenac	1.0	0.180	35.71
		10.0	0.098	65.00
		100.0	0.082	70.71
		0.1	0.232	17.14
3	Salicylanilide	1.0	0.194	30.71
		10.0	0.177	36.79
		100.0	0.151	46.07
		0.1	0.232	17.14




Figure 7. Graph of % Inhibition of Anti-Inflammatory Tests in Vitro

The results of the anti-inflammatory test for Diclofenac Sodium showed that the activity inhibited protein denaturation with a per cent inhibition of 25.00% at a concentration of 0.1 ppm and increased as the concentration increased to 1.0 ppm; 10.0 ppm and 100.0 ppm 35.71%, 65.00%, and 70.71%. The results of anti-inflammatory testing of the reaction compound showed activity to inhibit protein denaturation starting from a concentration of 1.0 ppm, namely with a % inhibition of 30.71%. The compound resulting from the reaction at a concentration of 0.1 ppm showed a % inhibition of 17.14%, which means that at this concentration, it did not have an anti-inflammatory effect; this is because

the % inhibition increased with increasing concentration so that at a concentration of 10.0 ppm it had a % inhibition of 36.79 % and at a concentration of 100.0 ppm it has an inhibition % of 46.07%. The results of in vitro anti-inflammatory testing show that the reaction compound has anti-inflammatory activity because it has a % inhibition value of >20%.

CONCLUSIONS

Salicylanilide compounds can be synthesized using sonochemical methods and produce the largest % yield in the 3rd sample at 5 hours at 11.23%. Salicylanilide compounds have anti-inflammatory activity at a concentration of 100.0 ppm with a % inhibition of 46.07%.

REFERENCES

- Almira, Diva, Windah Anugrah Subaidah, Agus Dwi Ananto, Rizqa Fersiyana Deccati, and Handa Muliasari. 2021. "In Vitro Concentration Optimization of Ethanol Extract from Makasar Fruit Seeds (*Brucea Javanica* L. Merr) as an Anti-Inflammatory Agent." *Jurnal Pijar Mipa* 16(5):595–99. doi: 10.29303/jpm.v16i5.2655.
- Draye, Micheline, Gregory Chatel, and Romain Duwald. 2020. "Ultrasound for Drug Synthesis: A Green Approach." *Pharmaceuticals* 13(2). doi: 10.3390/ph13020023.
- F. Dani Hendrata, Sabirin Matsjeh dan Iswahyudi. 2014. "Sintesis Metil 2-Asetoksibenzoat Dari Minyak Gandapura Dan Uji Aktivitasnya Sebagai Antiinflamasi the Synthesis of Methyl 2-Acetoxybenzoate From Wintergreen Oil and the Test As Antiinflammatory. 1–13."
- Farshori, Nida N., Mudasir R. Banday, Zeeshan Zahoor, and Abdul Rauf. 2010. "DCC/DMAP Mediated Esterification of Hydroxy and Non-Hydroxy Olefinic Fatty Acids with β -Sitosterol: In Vitro Antimicrobial Activity." *Chinese Chemical Letters* 21(6):646–50. doi: 10.1016/j.ccl.2010.01.003.
- Hendrata, F. Dani. 2014. "Sintesis Metil 2-Asetoksibenzoat Dari Minyak Gandapura Dan Uji Aktivitasnya Sebagai Antiinflamasi the Synthesis of Methyl 2-Acetoxybenzoate From Wintergreen Oil and the Test As Antiinflammatory." 1–13.
- Indriyanti, Erwin, and Masitoh Suryaning Prahasiwi. 2020. "Synthesis of Cinnamic Acid Based on Perkin Reaction Using Sonochemical Method and Its Potential as Photoprotective Agent." *JKPK (Jurnal Kimia Dan Pendidikan Kimia)* 5(1):54. doi: 10.20961/jkpk.v5i1.38136.
- Ma'mun. 2015. "Penyulingan Dan Analisis Beberapa Jenis Minyak Gandapura." *Buletin Penelitian Tanaman Rempah Dan Obat* 16(2):82–89.
- Manickam, Sivakumar, Daria Camilla Boffito, Erico M. M. Flores, Jean Marc Leveque, Rachel Pflieger, Bruno G. Pollet, and Muthupandian Ashokkumar. 2023. "Ultrasonics and Sonochemistry: Editors' Perspective." *Ultrasonics Sonochemistry* 99(July). doi: 10.1016/j.ultsonch.2023.106540.
- Nasution, Nur Azizah, Mala Nurilmala, and Asadatun Abdullah. 2019. "Seahorse Hydrolisate (*Hippocampus Kuda*) and Anti-Inflammatory Activity Test with Protein Denaturation Inhibition Method." *Jurnal Perikanan Universitas Gadjah Mada* 21(1):47. doi: 10.22146/jfs.43699.
- Of, Study, Analgesic Activity, O. F. Mucuna, Extract On, and Swiss Albino. 2015. "Study of Analgesic Activity of *Mucuna Pruriens*." 4(5):1124–33.
- Retnowati, Rurini, and Devina Azzuhro. 2013. "Sintesis N , N-Dietil-2-

- 1 Hidroksibenzamida Menggunakan Metil Salisilat Dari Minyak Gandapura.” *Natural B* 2(1):94–102.
- Ripa, Farhana Alam, Pritesh Ranjan Dash, and Md Omar Faruk. 2015. “CNS Depressant, Analgesic and Anti-Inflammatory Activities of Methanolic Seed Extract of Calamus Rotang Linn. Fruits in Rat.” *J Pharmacog Phytochem* 3(5):121–25.
- Sondhi, Sham M., Jaiveer Singh, Ashok Kumar, Hyder Jamal, and P. P. Gupta. 2009. “Synthesis of Amidine and Amide Derivatives and Their Evaluation for Anti-Inflammatory and Analgesic Activities.” *European Journal of Medicinal Chemistry* 44(3):1010–15. doi: 10.1016/j.ejmech.2008.06.029.
- Sulistyo, Rinda, Suratmo, and Rurini Retnowati. 2015. “Sintesis Salisilanilida Dari Komponen Utama Minyak Gandapura.” *Kimia Student Journal* 1(1):805–11.
- Whittaker, Joseph A., and Bernhard Vogler. 2008. “The in Vitro Anti-Denaturation Effects Induced by Natural Products and Non- Steroidal Compounds in Heat Treated (Immunogenic) Bovine Serum Albumin Is Proposed as a Screening Assay f ... The in Vitro Anti-Denaturation Effects Induced by Natural Products .” (October). doi: 10.1215/9780822388630-010.

Anti-inflammatory Activity of Salicylanilide Compounds from Gondopuro Oil as an In-Vitro

ORIGINALITY REPORT

19%	15%	0%	8%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	publikasiilmiah.unwahas.ac.id	9%
	Internet Source	
2	iosrphr.org	6%
	Internet Source	
3	Submitted to State Islamic University of Alauddin Makassar	5%
	Student Paper	

Exclude quotes	Off	Exclude matches	< 4%
Exclude bibliography	Off		