

BUKTI KORESPONDENSI
JURNAL INTERNASIONAL BEREPUTASI

Judul artikel : The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design

Jurnal : International Journal of Drug Delivery Technology (IJDDT)

Edisi : Tahun 2018, Vol 8, No.2 Halaman 110-115


Penerbit : Dr. Yashwant Research Labs Pvt. Ltd. (International Society for Science and Nature)

Penulis : Intan Martha Cahyani, Ebta Narasukma Anggraeny, Bektu Nugraheni, Christiana Retnaningsih, V Kristina Anani

Tabel Tahapan Publikasi Artikel

No	Tahapan Publikasi	Tanggal	Halaman
1	Manuscript Submission	25 Oktober 2017	Hal. 1
2	Manuscript Modification Notification	12 Januari 2018	Hal. 2
3	Follow-Up on manuscript modified request	12 Februari 2018	Hal. 2
4	Editorial Decision “Accepts accepted after minor modifications” and APC Payment request from IJDDT editorial	21 April 2018	Hal. 3
5	APC Payment Confirmation	3 Mei 2018	Hal. 3
6	Article Published Vol 8 issue 2 (25 Juni 2018) on website https://ijddt.com/volume8issue2/ https://impactfactor.org/PDF/IJDDT/8/IJDDT,Vol8,Issue2,Article12.pdf	25 Juni 2018	Hal. 4
7	IJDDT Published Information on Vol 8 issue 2 (25 Juni 2018)	4 Juli 2018	Hal. 5
8	Lampiran Bukti Korespondensi	-	Hal. 6 - 34

Pengusul


Dr. apt. Intan Martha Cahyani, M.Sc

1. Manuscript Submission

The screenshot shows a Gmail interface with a search bar containing "IJDDT". The email subject is "Re: 2017 Manuscript Submission (Perihal: Pengajuan Naskah 2017)". The sender is "IJDDT Journal <ijddtjournal@gmail.com>". The email body contains the following text:

Dear Author,
The manuscript is provided with number 10252017DDTC

On Sat, Oct 21, 2017 at 1:47 PM, intanmartha20@gmail.com <intanmartha20@gmail.com> wrote:

1. Name of Corresponding Author
intan cahyani

2. Email of Corresponding Author
intanmartha20@gmail.com

Institute of Affiliation of Corresponding Author
Sekolah Tinggi Ilmu Farmasi "Yayasan Pharmasi Semarang"

The screenshot shows a Gmail interface with a search bar containing "IJDDT". The email body contains the following text:

7. Keywords (2-10)
aqueous fraction, Clinacanthus nutans, microencapsulation, maltodextrin, Arabic gum

8. Complete manuscript in one file (word)

- [The-Optimization-of-Maltodextrin-and-Arabic-Gum-in-the-Microencapsulation-of-Aqueous-Fraction-of-Clinacanthus-nutans-Using-Simplex-Lattice-Design-FIX.doc](#)

9. Copyright Agreement form

- [Agreement-IJDDT.pdf](#)

--
Editor
International Journal of Drug Delivery Technology
ISSN: 0975-4415
website: www.ijddt.com

2. Manuscript Modification Notification

The screenshot shows a Gmail interface with a search bar containing "IJDDT". The left sidebar lists folders: Tulis, Kotak Masuk (1.804), Berbintang, Ditunda, Penting, Terkirim, Draf (43), Kategori, and Sosial (4). The main content area displays an email from "IJDDT Journal <ijddtjournal@gmail.com>" dated "12 Jan 2018, 20:29". The email body reads: "Dear Author, Please reply this mail with modified manuscript containing all inline references as superscripted numbers and match them with references written at the end. For better understanding please observe any other published manuscript." There is a button labeled "Terjemahkan ke Indonesia" and a "Show desktop" link at the bottom right.

3. Follow-Up on manuscript modified request

The screenshot shows a Gmail interface with a search bar containing "IJDDT". The left sidebar lists folders: Tulis, Kotak Masuk (1.804), Berbintang, Ditunda, Penting, Terkirim, Draf (43), Kategori, Sosial (4), Info Terbaru (1.296), Forum (1), Promosi (137), and Selengkapnya. The main content area displays an email from "Intan Martha Cahyani <intanmartha20@gmail.com>" dated "12 Feb 2018, 10:21". The email body reads: "Dear, International Journal of Drug Delivery Technology I attach improvements manuscript according to provisions that have been submitted". Below the text is a preview of an attachment titled "The Optimization ...". The email is followed by a response from "IJDDT Journal <ijddtjournal@gmail.com>" dated "21 Apr 2018, 10:40". There is a "Show desktop" link at the bottom right.

4. Editorial Decision “Accepts accepted after minor modifications” and APC Payment request from IJDDT editorial

The screenshot shows a Gmail interface with a search bar containing "IJDDT". The email is from "IJDDT Journal <ijddtjournal@gmail.com>" dated "21 Apr 2018, 10.40". The subject is "Dear author". The body text reads: "Your manuscript number " 10252017DDTC " bearing with title " The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of Clinacanthus nutans Using Simplex Lattice Design " has been provisionally **accepted after minor modifications**. It may be published in the upcoming issue of the journal subject to deposition of Open Access Fee USD150. The authors are required to use the **paypal link** available at this link www.ijddt.com/fee. Other modes are not acceptable. **Note:** The authors may **reply this mail** with modified manuscript containing minor typography errors if left coincidentally during submission. The errors must be highlighted clearly in **red color font**. The modifications are subject to approval of editorial board. Any typological/coincidental error left after this stage will not be considered. In case the modifications are not minor, then authors may be asked to withdraw and resubmit manuscript. **Important: After making a transaction, please confirm transaction by providing transaction details as a reply to this mail only.**

5. APC Payment Confirmation

The screenshot shows a Gmail interface with a search bar containing "IJDDT". The email is from "Intan Martha Cahyani <intanmartha20@gmail.com>" dated "3 Mei 2018, 17.45". The subject is "Dear IJDDT Journal,". The body text reads: "My manuscript number "10252017DDTC" bearing with title "The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of Clinacanthus nutans Using Simplex Lattice Design" Have been paid \$ 150.00 USD to Yashwant. (yash99yash@gmail.com) by PayPal ID transaction : 4XK45539LF129904A On May 3, 2018 15:08:28 GMT + 07:00. Please confirm if IJDDT Journal has received our payment We hope our article will be published in IJDDT Journal Thank you Regards

6. Article Published Vol 8 issue 2 (25 Juni 2018) on website

The screenshot shows the homepage of the International Journal of Drug Delivery Technology. The header includes the journal's logo, name, ISSN (0975 4415), and the text 'Peer Review Journal'. A navigation menu contains links for ARCHIVES, HOME, INSTRUCTIONS, EDITORIAL PROCESS, CONTACT, and SUBMIT MANUSCRIPT. A central banner displays 'Volume 8, Issue 2; April – June 2018'. Below this, two article titles are listed: '1. Partition Coefficient and Glutathione Penetration of Topical Antiaging: Preformulation Study' by Fitri Nugrahaeni, Dewi Melani Hariyadi, and Noorma Rosita; and '2. Formulation and Evaluation of Microemulsion Based Topical Gel of Carbamazepine' by Padhiar J S, Shirolkar S V, and Deshkar S S. A 'UGC approved Journal' badge is visible on the right. On the left, there are logos for 'Q3 Pharmaceutical Science' (best quartile) and 'Crossref'. The browser address bar shows 'https://ijddt.com/volume8issue2/'.

This screenshot shows the abstract page for the article '12. The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design' by Intan Martha Cahyani, Ebta Narasukma Anggraeny, Bekti Nugraheni, Christiana Retnaningsih, and V Kristina Ananingsih. The abstract text describes the microencapsulation of the aqueous fraction of *Clinacanthus nutans* leaf extracts using maltodextrin and Arabic gum. The study aimed to optimize the formulation using a simplex lattice design. The results showed that the optimum ratio of maltodextrin and Arabic gum was 0.806:0.194, with a moisture content of 1.49%, a flow rate of 4.375 g/s, and an antioxidant activity of 842,499 ppm. A 'View and Download Full Article' button is located at the bottom of the abstract. The footer of the page reads 'International Society for Science and Nature'. The browser address bar shows 'https://ijddt.com/volume8issue2/'.

7. IJDDT Published Information on Vol 8 issue 2

The screenshot shows a Gmail inbox on a web browser. The browser address bar displays the URL: <https://mail.google.com/mail/u/0/#search/IJDDT/FMfcgxmXJlpQMDhKNcwWPsdvwMfSgrT>. The Gmail interface includes a search bar with the text "IJDDT" and a left sidebar with various folders and categories. The main content area shows two email messages:

- Message 1:** From IJDDT Journal <ijddtjournal@gmail.com> to "saya" (me), dated 4 Jul 2018, 13:31. The subject is "Manuscript will be published in upcoming issue". It includes a "Terjemahkan ke Indonesia" button and a "Setelan" (Settings) dropdown menu.
- Message 2:** From Intan Martha Cahyani, dated 4 Jul 2018, 13:35. The subject is "Please find the attachment". It also includes a "Terjemahkan ke Indonesia" button.

The left sidebar shows the following categories and counts:

- Tulis
- Kotak Masuk: 1.804
- Berbintang
- Ditunda
- Penting
- Terkirim
- Draf: 43
- Kategori
- Sosial: 4
- Info Terbaru: 1.296
- Forum: 1
- Promosi: 137
- Selengkapnya

TRANSFER OF COPYRIGHT AGREEMENT

INTERNATIONAL JOURNAL OF DRUG DELIVERY TECHNOLOGY

Title The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design

Author(s) name(s): Intan Martha Cahyani¹⁾, Ebta Narasukma Anggraeny²⁾, Bekti Nugraheni³⁾, Christiana Retnaningsih⁴⁾, V. Kristina Ananingsih⁵⁾

Corresponding authors name address, affiliation and email Sekolah Tinggi Ilmu Farmasi “Yayasan Pharmasi Semarang” Jl. Letjend Sarwo Edie Wibowo KM. 1 Plamongansari Pucang Gading Semarang Jawa Tengah Indonesia 50193 email : intanmartha20@gmail.com

I warrant that the material / manuscript submitted to IJDDT is original and has not been and will be published, whole or in part, in print or electronic format, in any other journal.

Each author has participated sufficiently in the work to take public responsibility for appropriate portions of the content. The order of authorships on the byline is a joint decision of all the coauthors. In case the manuscript should be accepted for publication, I have the consent of each author to transfer and assign any and all right, title and interest, including copyright of the article referenced above.

I hereby assign and transfer copyright and all rights under it to the International Journal of Drug Delivery Technology.

Corresponding author Intan Martha Cahyani

Title The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design

Signature-----

The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design

, Intan Martha Cahyani¹⁾, Ebta Narasukma Anggraeny²⁾, Bektu Nugraheni³⁾, Christiana Retnaningsih⁴⁾,
V. Kristina Ananingsih⁵⁾

¹⁾²⁾³⁾ Sekolah Tinggi Ilmu Farmasi “Yayasan Pharmasi” Semarang

⁴⁾⁵⁾ Universitas Katolik Soegijapranata Semarang

intanmartha20@gmail.com

Abstracts

The aqueous fraction of *Clinacanthus nutans* leaf extracts contains flavonoids which known had antioxidative properties. To improve acceptability, this viscous and bitter aqueous fraction was microencapsulated using maltodextrin and Arabic gum. This research aims to discover the effectivity of maltodextrin and Arabic gum and the concentrations for optimum microencapsulation. Optimization design was done using Design Expert with simplex lattice design with ratios of 1:0; 0.75:0.25; 0.5:0.5; 0.25:0.75 and 0:1. The evaluations done to the results were microcapsule yield, moisture content, flow rate, and antioxidant activity. The optimum ratio of maltodextrin and Arabic gum was obtained at 0.806:0.194 with 1.49% moisture content, flow rate 4.375 g/s and antioxidant activity at the value of 842,499 ppm. The result of one-sample T-test showed that the prediction result of Design Expert did not differ significantly from the experiment result. From the data, it was concluded that the resulting equation was valid.

Keywords: aqueous fraction, *Clinacanthus nutans*, microencapsulation, maltodextrin, Arabic gum

Introduction

Currently, there are many research on plant antioxidants because the common synthetic ones such as BHA and BHT was suspected to be carcinogenic (Sayuti, 2015). *Clinacanthus nutans* is one of the plants commonly found in Indonesia and has potential as antioxidants. *Clinacanthus nutans* is a shrub commonly used as hedgerows in Indonesia. In traditional medicines, this plant is used as antidiabetics, antioxidants, antiinflammation, analgetics, antiviral, and also for wound treatment.

The leaves of *Clinacanthus nutans* contain alkaloids, triterpenoids/steroids, glycosides, tannins, saponins, and flavonoids. Based on research done by Akbar (2010), the isolates of *Clinacanthus nutans* methanolic leaf extracts contains six C-glycosides flavons, namely shaftoside, isomollupentin7-O- β -glukopyranoside, orientin, isoorientin, vitexin, and isovitexin. The flavonoids are known to acts as antioxidants by capturing free radicals. According to Anggraeny (2016), the aqueous fraction of *Clinacanthus nutans* leaf extracts had value of EC₅₀ 532,24 ppm, showing antioxidants properties.

The usage of *Clinacanthus nutans* leaf extracts fractions has several shortcomings, amongst them are its viscosity that gave difficulties in formulation and its astringent and bitter taste that made it hard to use orally. To overcome these shortcomings, microencapsulation can be used (Munin and Florence, 2011).

Microencapsulation is a process of microscopic encapsulation of drug particles with specific coat that result in better physical and chemical properties of those particles. Microencapsulation intends to protect sensitive components, reduce the loss of nutrients, and converts liquids to solids. The choice of microencapsulation methods depends on its applications and specific parameters, such as desired particle size, physicochemical properties of the core and the coating, release mechanism, process cost, etc. Freeze drying is one of drying method for encapsulation which had advantages in preserving the quality of the drying products, especially for heat-sensitive materials. Encapsulation can be attained as homogenous core in matrix solution which then was co-lyophilized, resulting in irregular shapes (Fang dan Bhandari, 2010). Freeze drying is a suitable method for encapsulation of antioxidants to preserve their properties because antioxidants are easily damaged by heat and light.

The choice of coating materials for microencapsulation of aqueous fraction of *Clinacanthus nutans* leaf extracts was based on Burin et.al. (2011), on the usage of combination of Arabic gum and maltodextrin for microencapsulation of grape anthocyanins and Mahdavi, et.al. on encapsulation of *Berberis vulgaris* extracts using Arabic gum and maltodextrin in 1:3 ratio. The combination of maltodextrin and Arabic gum as encapsulant is expected to form stable microcapsule and protect the aqueous fraction of *Clinacanthus nutans* because of the film-forming properties of Arabic gum and the ability of maltodextrin to protect microcapsules from oxidation. This research intended to find the optimum concentrations of microencapsulants of the aqueous fraction of *Clinacanthus nutans*.

Materials and Methods

Tools and Materials

Materials that were used on this research were *Clinacanthus nutans* leaves, analytical grade ethanol 96% (Merck), analytical grade methanol (Merck), analytical grade n-hexane (Merck), analytical grade ethyl acetate (Merck), DPPH, maltodextrin (*Dextrose Equivalent* 10,6), Arabic gum (*Tic Gums*), and distilled water. Tools used were separation funnels, rotary evaporator (*Buchi Rotavapor R-200*), freezer, freeze dryer (*Thermo Scientific Powerdry LL 1500*), moisture analyzer (*Mettler Toledo HE53*), flow funnels, ovens, UV-visible spectrophotometer (*Shimadzu*), SEM (*Scanning Electron Microscopy*).

Methods

Extraction dan Fractionation of *Clinacanthus nutans* Leaves Two hundred grams of *Clinacanthus nutans* leaf powder was macerated with two liters of ethanol 96% for two days. Diluted extracts was separated from the solvent using rotary vacuum evaporator at 40°C. The extract then gradually fractionated using liquid-liquid partition in separation funnels with water, ethyl acetate, and n-hexane as solvents. The water fraction then was evaporated using rotary vacuum evaporator at 80°C.

Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Leaf Extracts

Maltodextrin and Arabic gum in various ratio (see Table 1) were suspended in distilled water and the to the mixture the aqueous fraction of *Clinacanthus nutans* leaf extracts was added. The resulting mixture was frozen for 24 hours and then dried using freeze-dried in the temperature -100°C. The dry samples were pulverized and then were sifted using 24 mesh sieve. The resulting microcapsules were kept in tightly closed container and protected from light.

Table 1 Microcapsule Formulation of Aqueous Fraction *Clinacanthus nutans* Leaf Extracts

Materials	F1	F2	F3	F4	F5
Aqueous fraction (%)	4	4	4	4	4
Arabic gum	1	0,75	0,50	0,25	0
Maltodextrin	0	0,25	0,50	0,75	1

Note:

Maltodextrin and Arabic gum one part = 25 % and zero part = 15% each

The Evaluation of Microcapsule Characteristics

Microcapsule yield The yield was calculated by comparing the weight of microcapsule to the total active and coating materials (Ahn, et.al. 2007). The percentage of the microcapsule yield was calculated using Equation 1:

$$\text{Yield} = \frac{W_t - W_o}{W_o} \times 100\% \dots\dots\dots (1)$$

Note: Wt: initial weight (g)

Wo: end weight (g)

Moisture content. Moisture content was measured by weighing 0.5 grams of microcapsules and measuring them in the moisture meter. The start button was pressed and then the resulting numbers were notdn when the notification sounds were heard.

Solubility. Solubility were evaluated by solving one gram of microcapsules in 25 ml distilled water. The resulting solutions were filtered using Whatman paper No. 42. The filter papers and residues were dried in oven at 105°C for three hours and then were cooled down and weighed. The solubility percentage then was calculated using equations 2 and 3.

$$\text{Solubility percentage} = 100\% - \text{residue percentage} \dots\dots\dots (2)$$

$$\text{Residue percentage} = \frac{\text{Weight of filter paper and residue} - \text{weight of filter paper}}{\text{weight of sample}} \times 100\% \dots\dots\dots (3)$$

Flow Rate Flow rate was measured by weighing 100 g microcapsules, and then inputting those microcapsules into a closed-end funnel. The cover at the end of the funnel then was opened ant then the microcapsules was let to flow until there weren't any microcapsules remaining in the funnel. The flow time was recorded as the time the microcapsules needed to flow from the time the cover was opened until all the granules flowed out.

Antioxidant Activity Assay with DPPH Method Antioxidant activity was evaluated by solving 250 mg microcapsule in 50 ml of methanols and diluting the solution to 1000,1500, 2000, 2500 and 3000 ppm. From those solutions, 2 ml were taken and 4 ml DPPH solution

was added. The mixtures then were incubated at room temperature in dark condition for 30 minutes. The lowering of absorbancy was measured with spectrophotometer at 517 nm. The negative controls were made without any samples. From the resulted absorbancy, percentage of inhibition and EC₅₀ value were calculated. Percentage of inhibition was calculated using Equation 4.

$$\text{Percentage of inhibition} = \frac{\text{Negative control absorbancy} - \text{Sample absorbancy}}{\text{Negative control absorbancy}} \times 100\% \dots\dots\dots (4)$$

Scanning Electron Microscopy The optimized microcapsules was analyzed using SEM to examine particle size of the resulting microcapsules. The analysis was done by attaching the microcapsules to the holder using *dotile* then inputting the microcapsules into the vacuum evaporator. At certain degree of vacuum, the holder was set ablaze to make the gold vapor coat the tested material. The holder then was put into the instrument and the microcapsules were examined.

RESULTS AND DISCUSSION

The concentration range of maltodextrin and Arabic gum which can be used as encapsulant are were v from the total mixture. The ratio of aqueous fraction to the encapsulants was 1:10. The microencapsulation results can be seen at Figure 1.




Figure 1 Microcapsules of aqueous fraction of *Clinacanthus nutans* leaf extract

The result of physical characteristics and antioxidant test can be seen at table 2.

Tabel 2. result of physical characteristics and antioxidant test

Test	Formula				
	I	II	III	IV	V
Microcapsule yield (%)	28,15 ± 0,49	34,20 ± 1,38	53,48 ± 1,66	59,07 ± 1,47	40,05 ± 1,92
Moisture content (%)	1,32 ± 0,12	1,34 ± 0,06	1,67 ± 0,13	1,87 ± 0,04	2,06 ± 0,13
Solubility (%)	97,97 ± 0,52	97,16 ± 0,47	95,90 ± 0,35	95,84 ± 0,43	95,17 ± 0,22
Flow rate (g/s)	4,46 ± 0,26	4,00 ± 0,13	3,89 ± 0,19	3,90 ± 0,25	3,89 ± 0,19
Antioxidant activity (ppm)	919,98 ± 80,48	886,87 ± 56,88	703,37 ± 23,70	± 981,78 ± 96,21	± 919,88 ± 12,54

Note: The result above were the mean of four replications

Microcapsule yield was within the range of 28,15 ± 0,49 % to 59,07 ± 1,47%. Freeze-drying works by freezing the material and reducing the pressure around it and adding sufficient heat to enable the direct sublimation of ice (Fang dan Bhandari, 2010) therefore there are no other material other than water undergoes sublimation, even if the yield is less than 100%.

Moisture content tests were done to find the moisture content of the microcapsules. High moisture content result in damages of microcapsules and this damage influences the stability of microcapsules themselves. The moisture content of the aqueous fraction of *Clinacanthus nutans* leaf extracts microcapsules were found to be relatively low, within the range of 1.32 ± 0.12% to. 2,06 ± 0,13%. This low moisture content also can improve the flow rate of the granules. According to *Design Expert version 10.0 Trial* analysis results, it was shown that maltodextrin (M) and Arabic gum (G) each contributes significantly to the moisture content. On the other hand, the M-G interaction affected significantly in lowering moisture content. It was proven from the prob>F value smaller than 0.05 (>0.0001) according to Equation 6. The addition of Arabic gum can increase moisture content because Arabic gum contains a larger amount of shorter hydrophilic groups, therefore it easily binds water molecules in the air. (Mahdavi, 2016).

$$Y = 1,28 (M) + 2,08 (G) - 0,22 (MG) \dots \dots \dots (5)$$

Solubility test was done to find the capability of microcapsules to dissolve in water. The number of solubility value of the aqueous fraction of *Clinacanthus nutans* leaf extracts microcapsules was within the range of $95.17 \pm 0.22 \%$ to $97.97 \pm 0,52\%$. The solubility was affected by the moisture content, in which the low moisture content enable the microcapsule to disperse easily in water. According to the analysis results of *Design Experts version 10.0 Trial*, maltodextrin (M), Arabic gum (G), and MG interaction give significant effect with the prob>F value smaller than 0.05 (<0.0001) according to equation 7.

$$Y = 98.00(M) + 95,24(G) - 1,69(MG) \dots \dots \dots (6)$$

According to equation 7, it can be seen that the solubility will increase along the increase of the amount of maltodextrin dan Arabic gum. It was shown by the positive constants value. It can be seen from the equation that the addition of maltodextrin affected the increase of solubility since maltodextrin itself has high solubility (Rowe et.al., 2009).

Flow Rate Test was done to find the capability of certain amount of material to flow in certain time. High flow rate indicates that the material has good capability of flowing. Free-flowing microcapsules are highly desired because it makes measurement easier and ensures the homogeneity of composition and weight is all packages. The average flow rate of the resulting microcapsules was 4.03 g/s. According to Siregar (2008), a preparation can be said having good flow rate if it's flow rate is less than ten seconds. According to analysis results of *Design Experts version 10.0 Trial*, maltodextrin (M), Arabic gum (G), and MG interaction give significant effects , with prob>F less than 0.05 (<0.0011) according to Equation 8.

$$Y = 4,42(M) + 3,92(G) - 1,15(MG) \dots \dots \dots (7)$$

According to Equation 8, the value of flow rate wilin increase along the increase the maltodextrin because maltodextrin is a free-flowing material (Rowe et.al., 2009).

Antioxidant activity test. The test was done quantitatively using DPPH method. According to Molyneux (2004), the principle of this method is the measurement of the inhibition synthetic free radicals DPPH in polar organic solvents such as methanol by donating a hydrogen atom to stabilize purple DPPH radical to become yellow DPPH-H. The result then was compared to the antioxidant activity of the aqueous fraction of *Clinacanthus nutans* leaf extract. Accorning to the test, the EC₅₀ value of the microcapsules was within the range of 703.37 ± 23.70 to 919.98 ± 80.48 ppm while the antioxidant activity of aqueous fraction of *Clinacanthus nutans* leaf extract according to Nugraheni et.al. (2017) was 532.24 ppm. This reduction of antioxidant activity probably caused by the possibility of not all the fraction had been encapsulated. Although there was a reduction of antioxidant activity, the microcapsule can give protection to the aqueous fraction of *Clinacanthus nutans* leaf extract during the storage begause the active components are coated so their contact with air and light is reduced. According to the analysis result of *Design Experts version 10.0*, neither maltodextrin nor Arabic gum did not give significant effect to the antioxidant activity with the prob>F value larger than 0.05 (0.1331) which means it was not significant. The relating equation can be seen at Equation 9

$$Y = 921.19(M) + 959.08(G) - 462.09(MG) \dots \dots \dots (8)$$

The evaluation profile of the aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules according to the Simplex Lattice Design can be seen at Figure 2.




Figure 2. Profile of microcapsules of aqueous fraction of *Clinacanthus nutans* leaf extract according to the Simplex Lattice Design with (a) flow rate (b) moisture content (MC) (c) solubility (d) antioxidant activity.

The optimization process was done using *Design Experts version 10.0 Trial*. Then it would process all of the response variable according to the chosen criteria and give the solution formula as the chosen encapsulant formula. The formula which was chosen was the formula with desirability value namely the formula with 0.806 parts maltodextrin, 0.194 part Arabc gum. The formula had 4.141 g/s flow rate, 1.399 % moisture content, 97.203% solubility, and EC₅₀ 856.375 ppm. The resulting optimum formula can be seen at Figure 3.

Constraints						
Name	Goal	Lower Limit	Upper Limit	Lower Weight	Upper Weight	Importance
A:Maltodextrin	is in range	0	1	1	1	3
B:Gum Arab	is in range	0	1	1	1	3
Kecepatan Alir	maximize	3.57	4.76	0.1	1	3
MC	minimize	1.19	2.21	0.1	1	4
Kelarutan	maximize	94.89	98.53	0.1	1	5
Aktivitas Antioi	is target = 669	669.22	1114.26	0.1	1	5

Solutions							
Number	Maltodextrin	Gum Arab	Kecepatan Alir	MC	Kelarutan	Aktivitas Anti	Desirability
1	<u>0.806</u>	<u>0.194</u>	<u>4.141</u>	<u>1.399</u>	<u>97.203</u>	<u>856.375</u>	<u>0.786</u>

Figure 3 Formula optimum *Design Experts*

The resulting optimum formula then was validated to find whether the theoretical formula equation created by *Design Expert* would match the result of the experimental one. The theoretical result was compared to the experimental one with *one sample T-test* to find the validity of the optimization equation. The optimized formula for the microcapsules can be seen at Figure 4.



Figure 4. The Optimized microcapsules

Table 3. The experiment result compared to the theoretical value

Response	Actual value	Prediction value	Significance	Conclusion
Flow rate	4.375 g/s	4,141 g/s	0,088	Did not differ significantly
Moisture content	1.495 %	1,39 %	0,007	Differed significantly
Solubility	97.47 %	97,203 %	0,572	Did not differ significantly
EC ₅₀	842.499 ppm	856,371 ppm	0,010	Differed significantly

When the experiment results of each test parameters were compared to the theoretical ones, it was found that in the parameter of flow rate and solubility, it had significance larger than 0.05. It was then concluded that it did not differ significantly and the optimization equation was valid and could be used to predict the results.

SEM analysis then was done to the optimized microcapsules. This analysis intended to find the shape and the morphology of the microcapsules. The shapes and morphology can be seen at Figure 5. The microcapsules were not spherical, their surface was uneven, and they were porous. These shape was caused by the mixing and grinding process which could break the microcapsules to smaller particles with unspherical shapes. The uneven surface was caused by the imperfect process of polymer crosslinking.




Figure 5. The micrograph of aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules (a) 100x magnification (b) 1000x magnification (c) 2000x magnification

Conclusion

Based on the experiment result, it could be concluded that aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules could be made with the ratio of 0.804 part maltodextrin and 0.194 part Arabic gum. The resulting microcapsules had 1.49% moisture content; 97.47 % solubility; 4,375 g/s solubility; 842.499 ppm antioxidant activity.

Acknowledgement

The Ministry of Research, Technology, and Higher Education for the SKIM Inter-College Research Grant Number DIPA-042.06.1.401516/2017.

Bibliography

Ahn J.H., Kim Y.P., Seo E.M., Choi Y.K., and Kim H.S. 2007. Antioxidant Effect of Natural Plant Extracts on The Microencapsulated High Oleic Sunflower Oil. *Journal of Food Engine* 84:327-334.

Akbar, H, R. 2010. Isolasi dan Identifikasi Golongan Flavanoid Daun Dandang Gendis (*Clinacanthus nutans*) Berpotensi sebagai Antioksidan. *Skripsi*. Departemen Kimia Fakultas Matematika dan Ilmu Pengetahuan Alam IPB.

Burin V.M., P.N. Rossa, N.E. Ferreira-Lima, M.C.R. Hillmann, M.T.Boirdignon Luiz. 2011. Anthocyanins: optimisation of extraction from Cabernet Sauvignon grapes, microencapsulation and stability in soft drink, *Int. J. Food Sci. Technol.* 46 (1) :186–193.

Chelyn, J.L., Maizatul H.O., Nor, S.A.M.Y., Ramesh, R., Mohd, I.W., and Zakiah I. 2014. Analysis of Flavone C-Glycosides in the Leaves of *Clinacanthus nutans* (Burm. f.) Lindau by HPTLC and HPLC-UV/DAD. *The Scientific World Journal* 2014, Article ID 724267.

Fang, C., and Bhandari, B. 2010. Encapsulation Polyphenol. *Trends in Food Science & Technology* 21: 510 - 523

Jafari, Seid Mahdi., Katayoun Mahdavi Khazei., and Abbas Hemmati Kakhki. 2016. Microencapsulation of Saffron Petal Anthocyanins with cress seed Gum Compared with Arabic Gum through Freeze Drying. *Carbohydrate Polymers* 140: 20-25

Kembaren, R. B. 2012. Isolasi dan Nanoenkapsulasi Karotenoid Limbah Serat Kelapa Sawit. *Prosiding Seminar Ilmiah se-Eropa*

Mahdavi, S.A., Seid, M.J., Elham, A., and Danial D. 2016. Microencapsulation Optimization of Natural Anthocyanins with Maltodextrin, Gum Arabic and Gelatin. *International Journal of Biological Macromolecules* 85:379-385.

Munin, A and Florence, E. 2011. Encapsulation of Natural Polyphenolic Compounds; a Review. *France: Pharmaceutical Journal* ISSN 1999-4923. Hal 793-829

Molyneux, P. 2004. The Use of The Stable Free Radical Diphenylpicryl-hydrazil (DPPH) For Estimating Antioxidant Activity. *Songklanakarin Journal Science Technologi* 26 (2): 211-219.

Nugraheni, et al, 2017, Antioxidant Activity Test And Determination Of EC₅₀ Extract And Fraction Of Dandang Gendis Leaves (*Clinacanthus nutans*) By DPPH Method (1,1-Diphenyl-2-Picrylhydrazyl) In Vitro, Abstract, 2nd UMP-PIC & 8th ISCC (Pharmacy International Conference & Indonesian Society Of Cancer Chemoprevention): 68.

Rowe, R.C., Shesky, P.L., and Owen, S.C. (ed). 2009. *Handbook of Pharmaceutical Excipients* Edisi VI. London: The Pharmaceutical Press and The American Pharmacists Association.

Sayuti, K., and Rina Y. 2015. *Antioksidan, Alami dan Sintetik*. Padang: Andalas University Press

Siregar, and Charles J.P. 2008. *Teknologi Farmasi Sediaan Tablet : Dasar – Dasar Praktis*. Jakarta: Penerbit Buku Kedokteran, EGC.

The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design

Intan Martha Cahyani¹⁾, Ebita Narasukma Anggraeny²⁾, Bekti Nugraheni³⁾, Christiana Retnaningsih⁴⁾,
V. Kristina Ananingsih⁵⁾

¹⁾²⁾³⁾ Sekolah Tinggi Ilmu Farmasi “Yayasan Pharmasi” Semarang

⁴⁾⁵⁾ Universitas Katolik Soegijapranata Semarang

intanmartha20@gmail.com

ABSTRACT

The aqueous fraction of *Clinacanthus nutans* leaf extracts contains flavonoids which known had antioxidative properties. To improve acceptability, this viscous and bitter aqueous fraction was microencapsulated using maltodextrin and Arabic gum. This research aims to discover the effectivity of maltodextrin and Arabic gum and the concentrations for optimum microencapsulation. Optimization design was done using Design Expert with simplex lattice design with ratios of 1:0; 0.75:0.25; 0.5:0.5; 0.25:0.75 and 0:1. The evaluations done to the results were microcapsule yield, moisture content, flow rate, and antioxidant activity. The optimum ratio of maltodextrin and Arabic gum was obtained at 0.806:0.194 with 1.49% moisture content, flow rate 4.375 g/s and antioxidant activity at the value of 842,499 ppm. The result of one-sample T-test showed that the prediction result of Design Expert did not differ significantly from the experiment result. From the data, it was concluded that the resulting equation was valid.

Keywords: aqueous fraction, *Clinacanthus nutans*, microencapsulation, maltodextrin, Arabic gum

INTRODUCTION

Currently, there are many research on plant antioxidants because the common synthetic ones such as BHA and BHT was suspected to be carcinogenic¹⁰. *Clinacanthus nutans* is one of the plants commonly found in Indonesia and has potential as antioxidants. *Clinacanthus nutans* is a shrub commonly used as hedgerows in Indonesia. In traditional medicines, this plant is used as antidiabetics, antioxidants, antiinflammation, analgetics, antiviral, and also for wound treatment.

The leaves of *Clinacanthus nutans* contain alkaloids, triterpenoids/steroids, glycosides, tannins, saponins, and flavonoids. The isolates of *Clinacanthus nutans* methanolic leaf extracts contains six C-glycosides flavons, namely shaftoside, isomollupentin7-O- β -glukopyranoside, orientin, isoorientin, vitexin, and isovitexin². The flavonoids are known to acts as antioxidants by capturing free radicals. The aqueous fraction of *Clinacanthus nutans* leaf extracts had value of EC₅₀ 532,24 ppm, showing antioxidants properties⁸.

The usage of *Clinacanthus nutans* leaf extracts fractions has several shortcomings, amongst them are its viscosity that gave difficulties in formulation and its astringent and bitter taste that made it hard to use orally. To overcome these shortcomings, microencapsulation can be used⁶.

Microencapsulation is a process of microscopic encapsulation of drug particles with specific coat that result in better physical and chemical properties of those particles. Microencapsulation intends to protect sensitive components, reduce the loss of nutrients, and converts liquids to solids. The choice of microencapsulation methods depends on its applications and specific parameters, such as desired particle size, physicochemical properties of the core and the coating, release mechanism, process cost, etc. Freeze drying is one of drying method for encapsulation which had advantages in preserving the quality of the drying products, especially for heat-sensitive materials. Encapsulation can be attained as homogenous core in matrix solution which then was co-lyophilized, resulting in irregular shapes⁴. Freeze drying is a suitable method for encapsulation of antioxidants to preserve their properties because antioxidants are easily damaged by heat and light.

The choice of coating materials for microencapsulation of aqueous fraction of *Clinacanthus nutans* leaf extracts was based on the usage of combination of Arabic gum and maltodextrin for microencapsulation of grape anthocyanins³ and on encapsulation of *Berberis vulgaris* extracts using Arabic gum and maltodextrin in 1:3 ratio⁵. The combination of maltodextrin and Arabic gum as encapsulant is expected to form stable microcapsule and protect the aqueous fraction of *Clinacanthus nutans* because of the film-forming properties of Arabic gum and the ability of maltodextrin to protect microcapsules from oxidation. This research intended to find the optimum concentrations of microencapsulants of the aqueous fraction of *Clinacanthus nutans*.

MATERIALS AND METHODS

Tools and Materials

Materials that were used on this research were *Clinacanthus nutans* leaves, analytical grade ethanol 96% (Merck), analytical grade methanol (Merck), analytical grade n-hexane (Merck), analytical grade ethyl acetate (Merck), DPPH, maltodextrin (*Dextrose Equivalent* 10,6), Arabic gum (*Tic Gums*), and distilled water. Tools used were separation funnels, rotary evaporator (*Buchi Rotavapor R-200*), freezer, freeze dryer (*Thermo Scientific Powerdry LL 1500*), moisture analyzer (*Mettler Toledo HE53*), flow funnels, ovens, UV-visible spectrophotometer (*Shimadzu*), SEM (*Scanning Electron Microscopy*).

Methods

Extraction dan Fractionation of *Clinacanthus nutans* Leaves Two hundred grams of *Clinacanthus nutans* leaf powder was macerated with two liters of ethanol 96% for two days. Diluted extracts was separated from the solvent using rotary vacuum evaporator at 40°C. The extract then gradually fractionated using liquid-liquid partition in separation funnels with water, ethyl acetate, and n-hexane as solvents. The water fraction then was evaporated using rotary vacuum evaporator at 80°C.

Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Leaf Extracts Maltodextrin and Arabic gum in various ratio (see Table 1) were suspended in distilled water and the to the mixture the aqueous fraction of *Clinacanthus nutans* leaf extracts was added. The resulting mixture was frozen for 24 hours and then dried using freeze-dried in the temperature -100°C. The dry samples were pulverized and then were sifted using 24 mesh sieve. The resulting microcapsules were kept in tightly closed container and protected from light.

Table 1: Microcapsule Formulation of Aqueous Fraction *Clinacanthus nutans* Leaf Extracts

Materials	F1	F2	F3	F4	F5
Aqueous fraction (%)	4	4	4	4	4
Arabic gum	1	0,75	0,50	0,25	0
Maltodextrin	0	0,25	0,50	0,75	1

Note: Maltodextrin and Arabic gum one part = 25 % and zero part = 15% each

The Evaluation of Microcapsule Characteristics

Microcapsule yield The yield was calculated by comparing the weight of microcapsule to the total active and coating materials¹. The percentage of the microcapsule yield was calculated using Equation 1:

$$\text{Yield} = \frac{W_t - W_o}{W_o} \times 100\% \dots\dots\dots(1)$$

Note: Wt: initial weight (g)

W_o: end weight (g)

Moisture content. Moisture content was measured by weighing 0.5 grams of microcapsules and measuring them in the moisture meter. The start button was pressed and then the resulting numbers were noted when the notification sounds were heard.

Solubility. Solubility were evaluated by solving one gram of microcapsules in 25 ml distilled water. The resulting solutions were filtered using Whatman paper No. 42. The filter papers and residues were dried in oven at 105°C for three hours and then were cooled down and weighed. The solubility percentage then was calculated using equations 2 and 3.

$$\text{Solubility percentage} = 100\% - \text{residue percentage} \dots\dots\dots(2)$$

$$\text{Residue percentage} = \frac{\text{Weight of filter paper and residue} - \text{weight of filter paper}}{\text{weight of sample}} \times 100\% \dots\dots\dots (3)$$

Flow Rate Flow rate was measured by weighing 100 g microcapsules, and then inputting those microcapsules into a closed-end funnel. The cover at the end of the funnel then was opened ant then the microcapsules was let to flow until there weren't any microcapsules remaining in the funnel. The flow time was recorded as the time the microcapsules needed to flow from the time the cover was opened until all the granules flowed out.

Antioxidant Activity Assay with DPPH Method Antioxidant activity was evaluated by solving 250 mg microcapsule in 50 ml of methanols and diluting the solution to 1000,1500, 2000, 2500 and 3000 ppm. From those solutions, 2 ml were taken and 4 ml DPPH solution was added. The mixtures then were incubated at room temperature in dark condition for 30

minutes. The lowering of absorbancy was measured with spectrophotometer at 517 nm. The negative controls were made without any samples. From the resulted absorbancy, percentage of inhibition and EC₅₀ value were calculated. Percentage of inhibition was calculated using Equation 4.

$$\text{Percentage of inhibition} = \frac{\text{Negative control absorbancy} - \text{Sample absorbancy}}{\text{Negative control absorbancy}} \times 100\% \dots\dots\dots(4)$$

Scanning Electron Microscopy The optimized microcapsules was analyzed using SEM to examine particle size of the resulting microcapsules. The analysis was done by attaching the microcapsules to the holder using *dotile* then inputting the microcapsules into the vacuum evaporator. At certain degree of vacuum, the holder was set ablaze to make the gold vapor coat the tested material. The holder then was put into the instrument and the microcapsules were examined.

RESULTS AND DISCUSSION

The concentration range of maltodextrin and Arabic gum which can be used as encapsulant are were v from the total mixture. The ratio of aqueous fraction to the encapsulants was 1:10. The microencapsulation results can be seen at Figure 1. The result of physical characteristics and antioxidant test can be seen at table 2.




Figure 1: Microcapsules of aqueous fraction of *Clinacanthus nutans* leaf extract

Tabel 2: Result of physical characteristics and antioxidant test

Test	Formula				
	I	II	III	IV	V
Microcapsule yield (%)	28,15 ± 0,49	34,20 ± 1,38	53,48 ± 1,66	59,07 ± 1,47	40,05 ± 1,92
Moisture content (%)	1,32 ± 0,12	1,34 ± 0,06	1,67 ± 0,13	1,87 ± 0,04	2,06 ± 0,13
Solubility (%)	97,97 ± 0,52	97,16 ± 0,47	95,90 ± 0,35	95,84 ± 0,43	95,17 ± 0,22
Flow rate (g/s)	4,46 ± 0,26	4,00 ± 0,13	3,89 ± 0,19	3,90 ± 0,25	3,89 ± 0,19
Antioxidant activity (ppm)	919,98 ± 80,48	886,87 ± 56,88	703,37 ± 23,70	± 981,78 ± 96,21	± 919,88 ± 12,54

Note: The result above were the mean of four replications

Microcapsule yield was within the range of 28,15 ± 0,49 % to 59,07 ± 1,47%. Freeze-drying works by freezing the material and reducing the pressure around it and adding sufficient heat to enable the direct sublimation of ice⁴. therefore there are no other material other than water undergoes sublimation, even if the yield is less than 100%.

Moisture content tests were done to find the moisture content of the microcapsules. High moisture content result in damages of microcapsules and this damage influences the stability of microcapsules themselves. The moisture content of the aqueous fraction of *Clinacanthus nutans* leaf extracts microcapsules were found to be relatively low, within the range of 1.32 ± 0.12% to. 2,06 ± 0,13%. This low moisture content also can improve the flow rate of the granules. According to *Design Expert version 10.0 Trial* analysis results, it was shown that maltodextrin (M) and Arabic gum (G) each contributes significantly to the moisture content. On the other hand, the M-G interaction affected significantly in lowering moisture content. It was proven from the prob>F value smaller than 0.05 (>0.0001) according to Equation 6. The addition of Arabic gum can increase moisture content because Arabic gum contains a larger amount of shorter hydrophilic groups, therefore it easily binds water molecules in the air. (Mahdavi, et.al., 2016).

$$Y = 1,28 (M) + 2,08 (G) - 0,22 (MG) \dots\dots\dots(5)$$

Solubility test was done to find the capability of microcapsules to dissolve in water. The number of solubility value of the aqueous fraction of *Clinacanthus nutans* leaf extracts microcapsules was within the range of 95.17 ± 0.22 % to 97.97 ± 0,52%. The solubility was affected by the moisture content, in which the low moisture content enable the microcapsule

to disperse easily in water. According to the analysis results of *Design Expert version 10.0 Trial*, maltodextrin (M), Arabic gum (G), and MG interaction give significant effect with the prob>F value smaller than 0.05 (<0.0001) according to equation 7.

$$Y = 98.00(M) + 95,24(G) - 1,69(MG) \dots \dots \dots (6)$$

According to equation 7, it can be seen that the solubility will increase along the increase of the amount of maltodextrin dan Arabic gum. It was shown by the positive constants value. It can be seen from the equation that the addition of maltodextrin affected the increase of solubility since maltodextrin itself has high solubility⁹.

Flow Rate Test was done to find the capability of certain amount of material to flow in certain time. High flow rate indicates that the material has good capability of flowing. Free-flowing microcapsules are highly desired because it makes measurement easier and ensures the homogeneity of composition and weight is all packages. The average flow rate of the resulting microcapsules was 4.03 g/s. A preparation can be said having good flow rate if it's flow rate is less than ten seconds¹¹. According to analysis results of *Design Expert version 10.0 Trial*, maltodextrin (M), Arabic gum (G), and MG interaction give significant effects , with prob>F less than 0.05 (<0.0011) according to Equation 8.

$$Y = 4,42(M) + 3,92(G) - 1,15(MG) \dots \dots \dots (7)$$

According to Equation 8, the value of flow rate wilin increase along the increase the maltodextrin because maltodextrin is a free-flowing material⁹.

Antioxidant activity test. The test was done quantitatively using DPPH method. The principle of this method is the measurement of the inhibition synthetic free radicals DPPH in polar organic solvents such as methanol by donating a hydrogen atom to stabilize purple DPPH radical to become yellow DPPH-H⁷. The result then was compared to the antioxidant activity of the aqueous fraction of *Clinacanthus nutans* leaf extract. According to the test, the EC₅₀ value of the microcapsules was within the range of 703.37 ± 23.70 to 919.98 ± 80.48 ppm while the antioxidant activity of aqueous fraction of *Clinacanthus nutans* leaf extract was 532.24 ppm⁸. This reduction of antioxidant activity probably caused by the possibility of not all the fraction had been encapsulated. Although there was a reduction of antioxidant activity, the microcapsule can give protection to the aqueous fraction of *Clinacanthus nutans* leaf extract during the storage because the active components are coated so their contact with air and light is reduced. According to the analysis result of *Design Expert version 10.0 Trial*, neither maltodextrin nor Arabic gum did not give significant effect to the antioxidant activity with the prob>F value larger than 0.05 (0.1331) which means it was not significant. The relating equation can be seen at Equation 9

$$Y = 921.19(M) + 959.08(G) - 462.09(MG) \dots \dots \dots (8)$$

The evaluation profile of the aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules according to the Simplex Lattice Design can be seen at Figure 2.




Figure 2: Profile of microcapsules of aqueous fraction of *Clinacanthus nutans* leaf extract according to the Simplex Lattice Design with (a) flow rate (b) moisture content (MC) (c) solubility (d) antioxidant activity.

The optimization process was done using *Design Expert version 10.0 Trial*. Then it would process all of the response variable according to the chosen criteria and give the solution formula as the chosen encapsulant formula. The formula which was chosen was the formula with desirability value namely the formula with 0.806 parts maltodextrin, 0.194 part Arabic gum. The formula had 4.141 g/s flow rate, 1.399 % moisture content, 97.203% solubility, and EC_{50} 856.375 ppm. The resulting optimum formula can be seen at Figure 3.

Constraints						
Name	Goal	Lower Limit	Upper Limit	Lower Weight	Upper Weight	Importance
A:Maltodextrin	is in range	0	1	1	1	3
B:Gum Arab	is in range	0	1	1	1	3
Kecepatan Alir	maximize	3.57	4.76	0.1	1	3
MC	minimize	1.19	2.21	0.1	1	4
Kelarutan	maximize	94.89	98.53	0.1	1	5
Aktivitas Antio	is target = 669	669.22	1114.26	0.1	1	5

Solutions							
Number	Maltodextrin	Gum Arab	Kecepatan Alir	MC	Kelarutan	Aktivitas Anti	Desirability
1	<u>0.806</u>	<u>0.194</u>	<u>4.141</u>	<u>1.399</u>	<u>97.203</u>	<u>856.375</u>	<u>0.786</u> Selected

Figure 3: Formula optimum *Design Experts*

The resulting optimum formula then was validated to find whether the theoretical formula equation created by *Design Expert* would match the result of the experimental one. The theoretical result was compared to the experimental one with *one sample T-test* to find the validity of the optimization equation. The optimized formula for the microcapsules can be seen at Figure 4.



Figure 4: The Optimized microcapsules

Table 3: The experiment result compared to the theoretical value

Response	Actual value	Prediction value	Significance	Conclusion
Flow rate	4.375 g/s	4,141 g/s	0,088	Did not differ significantly
Moisture content	1.495 %	1,39 %	0,007	Differed significantly
Solubility	97.47 %	97,203 %	0,572	Did not differ significantly
EC ₅₀	842.499 ppm	856,371 ppm	0,010	Differed significantly

When the experiment results of each test parameters were compared to the theoretical ones, it was found that in the parameter of flow rate and solubility, it had significance larger than 0.05. It was then concluded that it did not differ significantly and the optimization equation was valid and could be used to predict the results.

SEM analysis then was done to the optimized microcapsules. This analysis intended to find the shape and the morphology of the microcapsules. The shapes and morphology can be seen at Figure 5. The microcapsules were not spherical, their surface was uneven, and they were porous. These shape was caused by the mixing and grinding process which could break the microcapsules to smaller particles with unspherical shapes. The uneven surface was caused by the imperfect process of polymer crosslinking.




Figure 5: The micrograph of aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules (a) 100x magnification (b) 1000x magnification (c) 2000x magnification

CONCLUSION

Based on the experiment result, it could be concluded that aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules could be made with the ratio of 0.804 part maltodextrin and 0.194 part Arabic gum. The resulting microcapsules had 1.49% moisture content; 97.47 % solubility; 4,375 g/s solubility; 842.499 ppm antioxidant activity.

ACKNOWLEDGEMENT

The Ministry of Research, Technology, and Higher Education for the SKIM Inter-College Research Grant Number DIPA-042.06.1.401516/2017.

REFERENCES

1. Ahn, J.H., Kim, Y.P., Seo, E.M., Choi, Y.K., and Kim, H.S. 2007. Antioxidant Effect of Natural Plant Extracts on The Microencapsulated High Oleic Sunflower Oil. *Journal of Food Engine* **84**:327-334.
2. Akbar, H.R. 2010. Isolasi dan Identifikasi Golongan Flavanoid Daun Dandang Gendis (*Clinacanthus nutans*) Berpotensi sebagai Antioksidan. *Skripsi*. Bandung: Departemen Kimia Fakultas Matematika dan Ilmu Pengetahuan Alam IPB.
3. Burin, V.M., Rossa, P.N., Ferreira-Lima, N.E., Hillmann, M.C.R., and Boirdignon-Luiz, M.T. 2011. Anthocyanins: Optimisation of Extraction from Cabernet Sauvignon Grapes, Microencapsulation and Stability in Soft Drink. *International Journal of Food Science and Technology* **46** (1):186–193.
4. Fang, C., and Bhandari, B. 2010. Encapsulation Polyphenol. *Trends in Food Science and Technology* **21**: 510 - 523
5. Mahdavi, S.A., Seid, M.J., Elham, A., and Danial, D. 2016. Microencapsulation Optimization of Natural Anthocyanins with Maltodextrin, Gum Arabic and Gelatin. *International Journal of Biological Macromolecules* **85**:379-385.
6. Munin, A and Florence, E. 2011. Encapsulation of Natural Polyphenolic Compounds; a Review. *Pharmaceutical Journal ISSN 1999-4923* :793-829.
7. Molyneux, P. 2004. The Use of The Stable Free Radical Diphenylpicryl-hydrazil (DPPH) For Estimating Antioxidant Activity. *Songklanakarinn Journal Science Technology* **26** (2): 211-219.
8. Nugraheni, B., Anggraeny, E.N., Cahyani, I.M., Retnaningsih, C., and Ananingsih, V.K. 2017. Antioxidant Activity Test And Determination Of EC₅₀ Extract And Fraction Of Dandang Gendis Leaves (*Clinacanthus nutans*) by DPPH Method (1,1-Diphenyl-2-Picrylhydrazyl) In Vitro. *Abstract. 2nd UMP-PIC & 8th ISCC (Pharmacy International Conference & Indonesian Society Of Cancer Chemoprevention)*: 68.
9. Rowe, R.C., Shesky, P.L., and Owen, S.C. (ed). 2009. *Handbook of Pharmaceutical Excipients*. Edisi VI. London: The Pharmaceutical Press and The American Pharmacists Association.
10. Sayuti, K., and Rina, Y. 2015. *Antioksidan, Alami dan Sintetik*. Padang: Andalas University Press
11. Siregar, and Charles J.P. 2008. *Teknologi Farmasi Sediaan Tablet : Dasar – Dasar Praktis*. Jakarta: Penerbit Buku Kedokteran, EGC.

The Optimization of Maltodextrin and Arabic Gum in the Microencapsulation of Aqueous Fraction of *Clinacanthus nutans* Using Simplex Lattice Design

Intan Martha Cahyani¹, Ebta Narasukma Anggraeny¹, Beki Nugraheni¹, Christiana Retnaningsih², V Kristina Ananingsih²

¹*Sekolah Tinggi Ilmu Farmasi "Yayasan Pharmasi" Semarang, Indonesia*

²*Universitas Katolik Soegijapranata Semarang, Indonesia*

Received: 21st Oct, 17; Revised: 12th Feb, 18, Accepted: 18th Mar, 18; Available Online: 25th Jun, 2018

ABSTRACT

The aqueous fraction of *Clinacanthus nutans* leaf extracts contains flavonoids which known had antioxidative properties. To improve acceptability, this viscous and bitter aqueous fraction was microencapsulated using maltodextrin and Arabic gum. This research aims to discover the effectivity of maltodextrin and Arabic gum and the concentrations for optimum microencapsulation. Optimization design was done using Design Expert with simplex lattice design with ratios of 1:0; 0.75:0.25; 0.5:0.5; 0.25:0.75 and 0:1. The evaluations done to the results were microcapsule yield, moisture content, flow rate, and antioxidant activity. The optimum ratio of maltodextrin and Arabic gum was obtained at 0.806:0.194 with 1.49% moisture content, flow rate 4.375 g/s and antioxidant activity at the value of 842,499 ppm. The result of one-sample T-test showed that the prediction result of Design Expert did not differ significantly from the experiment result. From the data, it was concluded that the resulting equation was valid.

Keywords: aqueous fraction, *Clinacanthus nutans*, microencapsulation, maltodextrin, Arabic gum.

INTRODUCTION

Currently, there are many research on plant antioxidants because the common synthetic ones such as BHA and BHT was suspected to be carcinogenic¹⁰. *Clinacanthus nutans* is one of the plants commonly found in Indonesia and has potential as antioxidants. *Clinacanthus nutans* is a shrub commonly used as hedgerows in Indonesia. In traditional medicines, this plant is used as antidiabetics, antioxidants, antiinflammation, analgetics, antiviral, and also for wound treatment.

The leaves of *Clinacanthus nutans* contain alkaloids, triterpenoids/steroids, glycosides, tannins, saponins, and flavonoids. The isolates of *Clinacanthus nutans* methanolic leaf extracts contains six C-glycosides flavons, namely shaftoside, isomollupentin7-O-β-glukopyranoside, orientin, isoorientin, vitexin, and isovitexin². The flavonoids are known to acts as antioxidants by capturing free radicals. The aqueous fraction of *Clinacanthus nutans* leaf extracts had value of EC₅₀ 532,24 ppm, showing antioxidants properties⁸.

The usage of *Clinacanthus nutans* leaf extracts fractions has several shortcomings, amongst them are its viscosity that gave difficulties in formulation and its astringent and bitter taste that made it hard to use orally. To overcome these shortcomings, microencapsulation can be used⁶.

Microencapsulation is a process of microscopic encapsulation of drug particles with specific coat that result in better physical and chemical properties of those

particles. Microencapsulation intends to protect sensitive components, reduce the loss of nutritions, and converts liquids to solids. The choice of microencapsulation methods depends on its applications and specific parameters, such as desired particle size, physicochemical properties of the core and the coating, release mechanism, process cost, etc. Freeze drying is one of drying method for encapsulation which had advantages in preserving the quality of the drying products, especially for heat-sensitive materials. Encapsulation can be attained as homogenous core in matrix solution which then was co-lyophilized, resulting in irregular shapes⁴. Freeze drying is a suitable method for encapsulation of antioxidants to preserve their properties because antioxidants are easily damaged by heat and light.

The choice of coating materials for microencapsulation of aqueous fraction of *Clinacanthus nutans* leaf extracts was based on the usage of combination of Arabic gum and maltodextrin for microencapsulation of grape anthocyanins³ and on encapsulation of *Berberis vulgaris* extracts using Arabic gum and maltodextrin in 1:3 ratio⁵. The combination of maltodextrin and Arabic gum as encapsulant is expected to form stable microcapsule and protect the aqueous fraction of *Clinacanthus nutans* because of the film-forming properties of Arabic gum and the ability of maltodextrin to protect microcapsules from oxidation. This research intended to find the optimum

concentrations of microencapsulants of the aqueous fraction of *Clinacanthus nutans*.

MATERIALS AND METHODS

Tools and Materials

Materials that were used on this research were *Clinacanthus nutans* leaves, analytical grade ethanol 96% (Merck), analytical grade methanol (Merck), analytical grade n-hexane (Merck), analytical grade ethyl acetate (Merck), DPPH, maltodextrin (*Dextrose Equivalent* 10,6), Arabic gum (*Tic Gums*), and distilled water. Tools used were separation funnels, rotary evaporator (*Buchi Rotavapor R-200*), freezer, freeze dryer (*Thermo Scientific Powerdry LL 1500*), moisture analyzer (*Mettler Toledo HE53*), flow funnels, ovens, UV-visible spectrophotometer (*Shimadzu*), SEM (*Scanning Electron Microscopy*).

Methods

Extraction dan Fractionation of Clinacanthus nutans Leaves

Two hundred grams of *Clinacanthus nutans* leaf powder was macerated with two liters of ethanol 96% for two days. Diluted extracts was separated from the solvent using rotary vacuum evaporator at 40°C. The extract then gradually fractionated using liquid-liquid partition in separation funnels with water, ethyl acetate, and n-hexane as solvents. The water fraction then was evaporated using rotary vacuum evaporator at 80°C.

Microencapsulation of Aqueous Fraction of Clinacanthus nutans Leaf Extracts

Maltodextrin and Arabic gum in various ratio (see Table 1) were suspended in distilled water and the to the mixture the aqueous fraction of *Clinacanthus nutans* leaf extracts was added. The resulting mixture was frozen for 24 hours and then dried using freeze-dried in the temperature -100°C. The dry samples were pulverized and then were sifted using 24 mesh sieve. The resulting microcapsules were kept in tightly closed container and protected from light.

The Evaluation of Microcapsule Characteristics

Microcapsule yield

The yield was calculated by comparing the weight of microcapsule to the total active and coating materials¹. The percentage of the microcapsule yield was calculated using Equation 1:

$$\text{Yield} = \frac{W_t - W_o}{W_o} \times 100\% \quad \dots\dots\dots(1)$$

Wt: initial weight (g)
Wo: end weight (g)

Moisture content

Moisture content was measured by weighing 0.5 grams of microcapsules and measuring them in the moisture meter. The start button was pressed and then the resulting numbers were noted when the notification sounds were heard.

Solubility

Solubility were evaluated by solving one gram of microcapsules in 25 ml distilled water. The resulting solutions were filtered using Whatman paper No. 42. The

Table 1: Microcapsule Formulation of Aqueous Fraction *Clinacanthus nutans* Leaf Extracts.

Materials	F1	F2	F3	F4	F5
Aqueous fraction (%)	4	4	4	4	4
Arabic gum	1	0,75	0,50	0,25	0
Maltodextrin	0	0,25	0,50	0,75	1




Figure 1: Microcapsules of aqueous fraction of *Clinacanthus nutans* leaf extract.

filter papers and residues were dried in oven at 105°C for three hours and then were cooled down and weighed. The solubility percentage then was calculated using equations 2 and 3.

$$\text{Solubility percentage} = 100\% - \text{residue percentage} \quad \dots\dots\dots(2)$$

$$\text{Residue percentage} = \frac{\text{Weight of filter paper and residue} - \text{weight of filter paper}}{\text{weight of sample}} \times 100\% \quad \dots\dots\dots(3)$$

Flow Rate

Flow rate was measured by weighing 100 g microcapsules, and then inputting those microcapsules into a closed-end funnel. The cover at the end of the funnel then was opened and then the microcapsules was let to flow until there weren't any microcapsules remaining in the funnel. The flow time was recorded as the time the microcapsules needed to flow from the time the cover was opened until all the granules flowed out.

Antioxidant Activity Assay with DPPH Method

Antioxidant activity was evaluated by solving 250 mg microcapsule in 50 ml of methanols and diluting the solution to 1000,1500, 2000, 2500 and 3000 ppm. From those solutions, 2 ml were taken and 4 ml DPPH solution was added. The mixtures then were incubated at room temperature in dark condition for 30 minutes. The lowering of absorbancy was measured with spectrophotometer at 517 nm. The negative controls were made without any samples. From the resulted absorbancy, percentage of inhibition and EC₅₀ value were calculated. Percentage of inhibition was calculated using Equation 4.

Table 2: Result of physical characteristics and antioxidant test.

Test	Formula				
	I	II	III	IV	V
Microcapsule yield (%)	28,15 ± 0,49	34,20 ± 1,38	53,48 ± 1,66	59,07 ± 1,47	40,05 ± 1,92
Moisture content (%)	1,32 ± 0,12	1,34 ± 0,06	1,67 ± 0,13	1,87 ± 0,04	2,06 ± 0,13
Solubility (%)	97,97 ± 0,52	97,16 ± 0,47	95,90 ± 0,35	95,84 ± 0,43	95,17 ± 0,22
Flow rate (g/s)	4,46 ± 0,26	4,00 ± 0,13	3,89 ± 0,19	3,90 ± 0,25	3,89 ± 0,19
Antioxidant activity (ppm)	919,98 ± 80,48	886,87 ± 56,88	703,37 ± 23,70	981,78 ± 96,21	919,88 ± 12,54

Note: The result above were the mean of four replications

$$\text{Percentage of inhibition} = \frac{\text{Negative control absorbancy} - \text{Sample absorbancy}}{\text{Negative control absorbancy}} \times 100\%$$

.....(4)

Scanning Electron Microscopy

The optimized microcapsules was analyzed using SEM to examine particle size of the resulting microcapsules. The analysis was done by attaching the microcapsules to the holder using *dotile* then inputting the microcapsules into the vacuum evaporator. At certain degree of vacuum, the holder was set ablaze to make the gold vapor coat the tested material. The holder then was put into the instrument and the microcapsules were examined.

RESULTS AND DISCUSSION

The concentration range of maltodextrin and Arabic gum which can be used as encapsulant are were v from the total mixture. The ratio of aqueous fraction to the encapsulants was 1:10. The microencapsulation results can be seen at Figure 1. The result of physical characteristics and antioxidant test can be seen at table 2.

Microcapsule yield was within the range of 28,15 ± 0,49 % to 59,07 ± 1,47%. Freeze-drying works by freezing the material and reducing the pressure around it and adding sufficient heat to enable the direct sublimation of ice⁴. therefore there are no other material other than water undergoes sublimation, even if the yield is less than 100%.

Moisture content tests were done to find the moisture content of the microcapsules. High moisture content result in damages of microcapsules and this damage influences the stability of microcapsules themselves. The moisture content of the aqueous fraction of *Clinacanthus nutans* leaf extracts microcapsules were found to be relatively low, within the range of 1.32 ± 0.12% to. 2,06 ± 0,13%. This low moisture content also can improve the flow rate of the granules. According to *Design Expert version 10.0 Trial* analysis results, it was shown that maltodextrin (M) and Arabic gum (G) each contributes significantly to the moisture content. On the other hand, the M-G interaction affected significantly in lowering moisture content. It was proven from the prob>F value smaller than 0.05 (>0.0001) according to Equation 6. The addition of Arabic gum can increase moisture content because Arabic gum contains a larger amount of shorter hydrophilic groups, therefore it easily binds water molecules in the air. (Mahdavi, et.al., 2016).

$$Y = 1,28 (M) + 2,08 (G) - 0,22 (MG)$$

.....(5)

Solubility test was done to find the capability of microcapsules to dissolve in water. The number of solubility value of the aqueous fraction of *Clinacanthus nutans* leaf extracts microcapsules was within the range of 95.17 ± 0.22 % to 97.97 ± 0,52%. The solubility was affected by the moisture content, in which the low moisture content enable the microcapsule to disperse easily in water. According to the analysis results of *Design Expert version 10.0 Trial*, maltodextrin (M), Arabic gum (G), and MG interaction give significant effect with the prob>F value smaller than 0.05 (<0.0001) according to equation 7.

$$Y = 98,00 (M) + 95,24 (G) - 1,69 (MG)$$

.....(6)

According to equation 7, it can be seen that the solubility will increase along the increase of the amount of maltodextrin dan Arabic gum. It was shown by the positive constants value. It can be seen from the equation that the addition of maltodextrin affected the increase of solubility since maltodextrin itself has high solubility⁹.

Flow Rate Test was done to find the capability of certain amount of material to flow in certain time. High flow rate indicates that the material has good capability of flowing. Free-flowing microcapsules are highly desired because it makes measurement easier and ensures the homogeneity of composition and weight is all packages. The average flow rate of the resulting microcapsules was 4.03 g/s. A preparation can be said having good flow rate if it's flow rate is less than ten seconds¹¹. According to analysis results of *Design Expert version 10.0 Trial*, maltodextrin (M), Arabic gum (G), and MG interaction give significant effects , with prob>F less than 0.05 (<0.0011) according to Equation 8.

$$Y = 4,42 (M) + 3,92 (G) - 1,15 (MG)$$

.....(7)

According to Equation 8, the value of flow rate wilin increase along the increase the maltodextrin because maltodextrin is a free-flowing material⁹.

Antioxidant activity test. The test was done quantitatively using DPPH method. The principle of this method is the measurement of the inhibition synthetic free radicals DPPH in polar organic solvents such as methanol by donating a hydrogen atom to stabilize purple DPPH radical to become yellow DPPH-H⁷. The result then was compared to the antioxidant activity of the aqueous fraction of *Clinacanthus nutans* leaf extract. According to the test, the EC₅₀ value of the microcapsules was within the range of 703.37 ± 23.70 to 919.98 ± 80.48 ppm while the antioxidant activity of aqueous fraction of *Clinacanthus nutans* leaf extract was 532.24 ppm⁸. This reduction of




Figure 2: Profile of microcapsules of aqueous fraction of *Clinacanthus nutans* leaf extract according to the Simplex Lattice Design with (a) flow rate (b) moisture content (MC) (c) solubility (d) antioxidant activity.

Constraints						
Name	Goal	Lower Limit	Upper Limit	Lower Weight	Upper Weight	Importance
A:Maltodextrin	is in range	0	1	1	1	3
B:Gum Arab	is in range	0	1	1	1	3
Kecepatan Alir	maximize	3.57	4.76	0.1	1	3
MC	minimize	1.19	2.21	0.1	1	4
Kelarutan	maximize	94.89	98.53	0.1	1	5
Aktivitas Antio	is target = 669	669.22	1114.26	0.1	1	5

Solutions							
Number	Maltodextrin	Gum Arab	Kecepatan Alir	MC	Kelarutan	Aktivitas Anti	Desirability
1	<u>0.806</u>	<u>0.194</u>	<u>4.141</u>	<u>1.399</u>	<u>97.203</u>	<u>856.375</u>	<u>0.786</u> Selected

Figure 3: Formula optimum *Design Experts*.



Figure 4: The Optimized microcapsules

antioxidant activity probably caused by the possibility of not all the fraction had been encapsulated. Although there was a reduction of antioxidant activity, the microcapsule can give protection to the aqueous fraction of *Clinacanthus nutans* leaf extract during the storage because the active components are coated so their contact with air and light is reduced. According to the analysis result of *Design Expert version 10.0 Trial*, neither maltodextrin nor Arabic gum did not give significant effect to the antioxidant activity with the prob>F value larger than 0.05 (0.1331) which

Table 3: The experiment result compared to the theoretical value.

Response	Actual value	Prediction value	Significance	Conclusion
Flow rate	4.375 g/s	4,141 g/s	0,088	Did not differ significantly
Moisture content	1.495 %	1,39 %	0,007	Differed significantly
Solubility	97.47 %	97,203 %	0,572	Did not differ significantly
EC ₅₀	842.499 ppm	856,371 ppm	0,010	Differed significantly




Figure 5: The micrograph of aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules (a) 100x magnification (b) 1000x magnification (c) 2000x magnification

means it was not significant. The relating equation can be seen at Equation 9

$$Y = 921.19 (M) + 959.08 (G) - 462.09 (MG) \dots \dots \dots (8)$$

The evaluation profile of the aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules according to the Simplex Lattice Design can be seen at Figure 2.

The optimization process was done using *Design Expert version 10.0 Trial*. Then it would process all of the response variable according to the chosen criteria and give the solution formula as the chosen encapsulant formula. The formula which was chosen was the formula with desirability value namely the formula with 0.806 parts maltodextrin, 0.194 part Arabic gum. The formula had 4.141 g/s flow rate, 1.399 % moisture content, 97.203% solubility, and EC₅₀ 856.375 ppm. The resulting optimum formula can be seen at Figure 3.

The resulting optimum formula then was validated to find whether the theoretical formula equation created by *Design Expert* would match the result of the experimental one. The theoretical result was compared to the experimental one with *one sample T-test* to find the validity of the optimization equation. The optimized formula for the microcapsules can be seen at Figure 4.

When the experiment results of each test parameters were compared to the theoretical ones, it was found that in the parameter of flow rate and solubility, it had significance larger than 0.05. It was then concluded that it did not differ significantly and the optimization equation was valid and could be used to predict the results.

SEM analysis then was done to the optimized microcapsules. This analysis intended to find the shape and the morphology of the microcapsules. The shapes and morphology can be seen at Figure 5. The microcapsules were not spherical, their surface was uneven, and they were porous. These shape was caused by the mixing and grinding process which could break the microcapsules to smaller particles with unspherical shapes. The uneven surface was caused by the imperfect process of polymer crosslinking.

CONCLUSION

Based on the experiment result, it could be concluded that aqueous fraction of *Clinacanthus nutans* leaf extract microcapsules could be made with the ratio of 0.804 part maltodextrin and 0.194 part Arabic gum. The resulting microcapsules had 1.49% moisture content; 97.47 % solubility; 4,375 g/s solubility; 842.499 ppm antioxidant activity.

ACKNOWLEDGEMENT

The Ministry of Research, Technology, and Higher Education for the SKIM Inter-College Research Grant Number DIPA-042.06.1.401516/2017.

REFERENCES

- Ahn, J.H., Kim, Y.P., Seo, E.M., Choi, Y.K., and Kim, H.S. 2007. Antioxidant Effect of Natural Plant Extracts on The Microencapsulated High Oleic Sunflower Oil. *Journal of Food Engine* 84:327-334.
- Akbar, H.R. 2010. Isolasi dan Identifikasi Golongan Flavanoid Daun Dandang Gendis (*Clinacanthus nutans*) Berpotensi sebagai Antioksidan. *Skripsi*. Bandung: Departemen Kimia Fakultas Matematika dan Ilmu Pengetahuan Alam IPB.
- Burin, V.M., Rossa, P.N., Ferreira-Lima, N.E., Hillmann, M.C.R., and Boirdignon-Luiz, M.T. 2011. Anthocyanins: Optimisation of Extraction from Cabernet Sauvignon Grapes, Microencapsulation and Stability in Soft Drink. *International Journal of Food Science and Technology* 46 (1):186–193.
- Fang, C., and Bhandari, B. 2010. Encapsulation Polyphenol. *Trends in Food Science and Technology* 21: 510 - 523
- Mahdavi, S.A., Seid, M.J., Elham, A., and Danial, D. 2016. Microencapsulation Optimization of Natural Anthocyanins with Maltodextrin, Gum Arabic and Gelatin. *International Journal of Biological Macromolecules* 85:379-385.

6. Munin, A and Florence, E. 2011. Encapsulation of Natural Polyphenolic Compounds; a Review. *Pharmaceutical Journal* ISSN 1999-4923 :793-829.
7. Molyneux, P. 2004. The Use of The Stable Free Radical Diphenylpicryl-hydrazil (DPPH) For Estimating Antioxidant Activity. *Songklanakarinn Journal Science Technology* 26 (2): 211-219.
8. Nugraheni, B., Anggraeny, E.N., Cahyani, I.M., Retnaningsih, C., and Ananingsih, V.K. 2017. Antioxidant Activity Test And Determination Of EC₅₀ Extract And Fraction Of Dandang Gendis Leaves (*Clinacanthus nutans*) by DPPH Method (1,1-Diphenyl-2-Picrylhydrazyl) In Vitro. *Abstract*. 2nd UMP-PIC & 8th ISCC (Pharmacy International Conference & Indonesian Society Of Cancer Chemoprevention): 68.
9. Rowe, R.C., Shesky, P.L., and Owen, S.C. (ed). 2009. *Handbook of Pharmaceutical Excipients*. Edisi VI. London: The Pharmaceutical Press and The American Pharmacists Association.
10. Sayuti, K., and Rina, Y. 2015. *Antioksidan, Alami dan Sintetik*. Padang: Andalas University Press
11. Siregar, and Charles J.P. 2008. *Teknologi Farmasi Sediaan Tablet : Dasar – Dasar Praktis*. Jakarta: Penerbit Buku Kedokteran, EGC.